arXiv:1403.2749v1 [math.CO] 11 Mar 2014

Embedding multidimensional grids into optimal hypercubes

Zevi Miller * Dan Pritikin † I.H. Sudborough ‡

March 13, 2014

Abstract

Let G and H be graphs, with $|V(H)| \geq |V(G)|$, and $f : V(G) \to V(H)$ a one to one map of their vertices. Let $dilation(f) = max\{dist_H(f(x), f(y)) : xy \in E(G)\}$, where $dist_H(v, w)$ is the distance between vertices v and w of H. Now let $B(G, H) = min_f\{dilation(f)\}$, over all such maps f.

The parameter $B(G, H)$ is a generalization of the classic and well studied "bandwidth" of G, defined as $B(G, P(n))$, where $P(n)$ is the path on n points and $n = |V(G)|$. Let $[a_1 \times a_2 \times \cdots \times a_k]$ be the k-dimensional grid graph with integer values 1 through a_i in the i'th coordinate. In this paper, we study $B(G, H)$ in the case when $G = [a_1 \times a_2 \times \cdots \times a_k]$ and H is the hypercube Q_n of dimension $n = \lceil log_2(|V(G)|)\rceil$, the hypercube of smallest dimension having at least as many points as G. Our main result is that

$$B([a_1 \times a_2 \times \cdots \times a_k], Q_n) \leq 3k,$$

provided $a_i \geq 2^{22}$ for each $1 \leq i \leq k$. For such G, the bound $3k$ improves on the previous best upper bound $4k + O(1)$. Our methods include an application of Knuth's result on two-way rounding and of the existence of spanning regular cyclic caterpillars in the hypercube.

1 Introduction

In this paper we will usually follow standard graph theoretic terminology, as may be found for example in [28]. We let $P(t)$ stand for the path on t vertices. The cartesian product $G \times H$ of two graphs G and H is the graph with vertex set $V = \{(v, w) : v \in V(G), w \in V(H)\}$ and edge set $E = \{(v, w)(v', w') : \text{either } v = v' \text{ and } ww' \in E(H), \text{ or } vv' \in E(G) \text{ and } w = w'\}$. All logarithms are taken base 2.

*Dept. of Mathematics, Miami University, Oxford, OH 45056, USA, millerz@muohio.edu
†Dept. of Mathematics, Miami University, Oxford, OH 45056, USA, pritikd@muohio.edu.
‡Program in Computer Science, University of Texas-Dallas, Richardson, Texas

1.1 Background and main result

The analysis of how effectively one network can simulate another, and the resulting implications for optimal design of parallel computation networks, are important topics in graph theoretic aspects of computer science. One of the measures of the effectiveness of a simulation is the *dilation* of the corresponding map (or "embedding") of networks, defined as follows. Let G and H be two graphs and $f : V(G) \to V(H)$ a map from the vertices of G to those of H. As a convenience we typically write such a map as $f : G \to H$, with the meaning that it is a map from vertices to vertices. Similarly we sometimes write $|G|$ for $|V(G)|$. Apart from an exception indicated below in a review of previous research on our topic, we will suppose that $|V(G)| \leq |V(H)|$ and that f is one to one. Whether f is one to one or not, we let $dilation(f) = max\{dist_H(f(x), f(y)) : xy \in E(G)\}$, where $dist_H(v, w)$ is the distance between vertices v and w of H, defined as the minimum number of edges in any path of H joining v and w. Thus $dilation(f)$ is the maximum "stretch" experienced by any edge of G under the map f. Now define $B(G, H)$ to be $min_f\{dilation(f)\}$, over all such maps f. Note that $B(G, H)$ is a generalization of the classic and well studied "bandwidth" of G, defined as $B(G, P(n))$, where $n = |V(G)|$.

The study of $B(G, H)$ arises when each of G and H is a computation network, and the goal is to have H simulate a computation in G. A given map f indicates how the vertices of H play the roles of the vertices of G, and $dilation(f)$ is a measure of the communication delay in this roleplaying. A message between adjacent vertices x and y in G taking unit time would become a message between $f(x)$ and $f(y)$ in H taking time $dist_H(f(x), f(y))$, which in the worst case is $dilation(f)$ if a shortest path in H joining $f(x)$ and $f(y)$ for this message is used. Indeed the delay may be worse when one considers the full simulation, requiring in addition to f a *routing path* for each edge $xy \in E(G)$, namely, a path in H (not necessarily shortest) joining $f(x)$ and $f(y)$. So let the *edge congestion* of f be the maximum, over all edges $vw \in E(H)$, of the number of routing paths in H that contain vw. The edge congestion of f is then an additional contribution to the communication delay of the embedding f.

In this paper we obtain upper bounds on $B(G, H)$ when G is a multidimensional grid and H is the smallest hypercube having at least $|V(G)|$ vertices. To clarify, let $a_i \geq 2$, $1 \leq i \leq k$, be integers. The k-dimensional grid $G = [a_1 \times a_2 \times \cdots \times a_k]$ is the graph with vertex set $V(G) = \{x = (x_1, x_2, \ldots, x_k) : x_i \text{ an integer, } 1 \leq x_i \leq a_i\}$ and edge set $E(G) = \{xy : \sum_{i=1}^{k} |x_i - y_i| = 1\}$. So two vertices of G are joined by an edge precisely when they disagree in exactly one coordinate, and in that coordinate they differ by 1. It follows that for $x, y \in V(G)$ we have $dist_G(x, y) = \sum_{i=1}^{k} |x_i - y_i|$. One can also write G as the cartesian product of paths $G = P(a_1) \times P(a_2) \times \cdots \times P(a_k)$.

The *n-dimensional hypercube* Q_n is the n-dimensional grid $[2 \times 2 \times \cdots \times 2]$. We follow the traditional view whereby $V(Q_n)$ is the set of all strings of length n over the alphabet $\{0, 1\}$, where two such strings are joined by an edge if they disagree in exactly one coordinate. This departs in a trivial way from our notation above, where we would have required $1 \leq x_i \leq 2$. Clearly $|V(Q_n)| = 2^n$ and we let $Opt(G)$ be the smallest hypercube containing at least $|V(G)|$ vertices, so $Opt(G) = Q_t$ where $t = \lceil log_2(|V(G)|) \rceil$.

There is a substantial literature on the simulation of various networks by hypercubes and their related networks; the butterfly, shuffle exchange and DeBruijn graphs. See the books [21] and [25] for excellent expositions on these topics, both emphasizing bounds on dilation and

congestion in graph embeddings; where the first also includes routing and implementation of various algorithms while the second gives a unified approach to applying separator theorems for deriving such bounds. An early survey on embedding graphs into hypercubes [22] mentions necessary and sufficient conditions (originating in [16]) for a graph to be a subgraph of some hypercube, and also the fact that for the complete binary tree T_n on $2^n - 1$ vertices there is an embedding $f : T_n \to Q_n$ such that for every edge $xy \in E(T_n)$ we have $dist_{Q_n}(f(x), f(y)) = 1$ with the exception of a single edge where this distance is 2 [15]. In [6] it is shown how to embed any 2^n node bounded degree tree into Q_n with $O(1)$ dilation and $O(1)$ edge congestion, as n grows. In the same paper these results are extended to embedding bounded degree graphs with $O(1)$ separators. In [21] many-to-one maps of binary trees into hypercubes are considered, letting the *load* be the maximum number of tree nodes mapped onto a hypercube node. Using probabilistic methods and error correcting codes it is shown how to embed an M node binary tree in an N node hypercube with dilation 1 and load $O(\frac{M}{N} + log(N))$, and how to perform the same embedding with dilation $O(1)$ and load $O(\frac{M}{N} + 1)$.

Concerning the embedding of multidimensional grids into hypercubes, observe first that if $p_1, p_2, \ldots, p_r$ are positive integers summing to n, and $G = [P(2^{p_1}) \times P(2^{p_2}) \times \cdots \times P(2^{p_r})]$, then $Q_n = Opt(G)$ and Q_n contains G as a spanning subgraph. Thus $B(G, Opt(G)) = 1$ in this case. In fact one can show that $[a_1 \times a_2 \times \cdots \times a_k]$ is a subgraph of Q_n if and only if $n \geq \lceil log(a_1) \rceil + \lceil log(a_2) \rceil + \cdots + \lceil log(a_k) \rceil$; see Problem 3.20 in [21]. Answering a question posed in [22] about 2-dimensional grids $G = [a_1 \times a_2]$, it is shown in [9] and in [8] that $B(G, Opt(G)) \leq 2$. In [9] it is also shown for arbitrary multidimensional grids $G = [a_1 \times a_2 \times \cdots \times a_k]$ that $B(G, Opt(G)) \leq 4k + 1$. Independently it was shown in [20] that $B(G, Opt(G)) \leq 4k - 1$ for such G, this upper bound being realized by a parallel algorithm on the hypercube. Still for such G, it was shown in [5] that $B(G, Opt(G)) \leq k$, assuming quite involved and restrictive inequality constraints on the a_i. Returning to dimension 2, it was shown in [18] that determining whether a given graph G can be embedded in $Opt(G)$ with edge congestion 1 is NP-complete. Subsequently it was shown in [26] that any $G = [a_1 \times a_2]$ can be embedded in $Opt(G)$ with edge congestion at most 2 and dilation at most 3. Following up on a question posed in [22], the issue of many-to-one embeddings of 2 and 3 dimensional grids G into hypercubes was explored in [24]. For these results, let $Opt(G)/2^t$ denote the hypercube of dimension $\lceil log(|G|) \rceil - t$. If $f : G \to Opt(G)/2^t$ is a many-to-one map, then as above let the *load* of f be $\max\{|f^{-1}(z)| : z \in Opt(G)/2^t\}$. It was shown in [24] that for a 2-dimensional grid G there is a many-to-one map $f : G \to Opt(G)/2^t$ of dilation 1 and load at most $1 + 2^t$, and when G is 3-dimensional there is a map $f : G \to Opt(G)/2$ of dilation at most 2 and load at most 3, and a map $f : G \to Opt(G)/4$ of dilation at most 3 and load at most 5.

The main result of the present paper is that $B([a_1 \times a_2 \times \cdots \times a_k], Q_n) \leq 3k$, provided $a_i \geq 2^{22}$ for each $1 \leq i \leq k$. This improves on the $4k - 1$ bound above under this condition on the a_i. We construct a one to one map $H^k : G \to Opt(G)$ realizing $dilation(H^k) \leq 3k$ and having congestion $O(k)$. Our construction uses the technique of two way rounding and the existence of regular spanning cyclic caterpillars in the hypercube.

1.2 Some notation

We will need to consider multidimensional grids for which each factor (in the cartesian product) with one possible exception is a path $P(m)$, where m is a power of 2 and varies with the factor, as these will play the role of successive approximations to $Opt(G)$. Let $e_i = \lceil log_2(a_1 a_2 \cdots a_i) \rceil$ for $1 \leq i \leq k$, with $e_0 = 0$. Letting $p_i = e_i - e_{i-1}$ for $1 \leq i \leq k$, we let $Opt'(G) = P(2^{p_1}) \times P(2^{p_2}) \times \cdots \times P(2^{p_k})$. So $Opt'(G)$ is a spanning subgraph of $Opt(G)$. For any $1 \leq t \leq k$ let $\langle Y_t \rangle = P(2^{p_1}) \times P(2^{p_2}) \times \cdots \times P(2^{p_t})$, with these p_i. We let $Y_t = \langle Y_{t-1} \rangle \times P(l)$, where l is large enough. Thus Y_t is the t-dimensional grid $P(2^{e_1}) \times P(2^{e_2 - e_1}) \times \cdots \times P(2^{e_{t-1} - e_{t-2}}) \times P(l)$. The grids Y_2, Y_3,... will be the aforementioned successive approximations to $Opt(G)$. We will construct one to one maps $f_i : G \to Y_i$, $2 \leq i \leq k$. The final map f_k will satisfy $f_k(G) \subseteq Opt'(G) \subseteq Opt(G)$.

For any point x in a multidimensional grid, we let x_i be its i'th coordinate (as suggested above), and when $i \leq j$ we let $x_{i \to j}$ be the $(j - i + 1)$-tuple $(x_i, x_{i+1}, \ldots, x_j)$. So for example, let $f : G \to H$ be a map where G and H are both multidimensional grids and H is of dimension r. Let $f(x) = (b_1, b_2, \cdots, b_r) \in V(H)$ for some $x \in V(G)$. Then by our notation $f(x)_2 = b_2$, while $f(x)_{1 \to i} = (b_1, b_2, \cdots, b_i)$. We can also express $dilation(f)$ as $dilation(f) = \max \{\sum_{i=1}^{r} |f(x)_i - f(y)_i| : xy \in E(G)\}$.

For $1 \leq t < i$, a *t-level* of Y_i is any t-dimensional subgrid of Y_i obtained by fixing the last $i - t$ coordinates of points in Y_i. Recalling that $Y_i = \langle Y_{i-1} \rangle \times P(l)$, note that there are l pairwise disjoint $(i-1)$-levels of Y_i, each isomorphic to $\langle Y_{i-1} \rangle$, and we denote by Y_i^c the $(i-1)$-level all of whose points have last (that is, i'th) coordinate c, $1 \leq c \leq l$. We also let $Y_i^{(r)} = \bigcup_{c=1}^{r} Y_i^c$. For fixed i and j, $2 \leq i \leq k$ and $j \geq 1$, we denote by S_i^j the subgraph of Y_i induced by the vertices $\{(x_1, x_2, \ldots, x_{i-1}, y) \in Y_i : 1 \leq x_q \leq 2^{e_q - e_{q-1}}$ for $1 \leq q \leq i-1, 1 + (j-1)2^{e_i - e_{i-1}} \leq y \leq j2^{e_i - e_{i-1}}\}$, and call S_i^j an *i-section* of Y_i, more precisely, the j'th *i-section*. Thus we have $S_i^j \cong \langle Y_{i-1} \rangle \times P(2^{e_i - e_{i-1}}) \cong \langle Y_i \rangle$ for each j, and we let $S_i^{(r)} = \bigcup_{j=1}^{r} S_i^j$.

As an example, let $G = [3 \times 7 \times 5 \times 9]$. Then $e_1 = 2$, $e_2 = 5$, $e_3 = 7$, $\langle Y_2 \rangle = [4 \times 8]$, and $Y_2 = 4 \times l$ for large l. Further, $Y_2^3 = \{(x_1, 3) : 1 \leq x_1 \leq 4\}$, so Y_2^3 can be thought of as the third column of Y_2, and $Y_2^{(3)} = \cup_{j=1}^{3} Y_2^j$ is the graph induced by the union of the first three columns of Y_2. We have $S_2^j \cong \langle Y_2 \rangle = [4 \times 8]$ for each $j \geq 1$, and S_2^j is the $[4 \times 8]$ subgrid of Y_2 induced by columns $8(j-1)+1$ through $8j$. We have $\langle Y_3 \rangle = [4 \times 8 \times 4]$, $Y_3 = [4 \times 8 \times l]$ for large l, and $Y_3^c = \{(x_1, x_2, c) : 1 \leq x_1 \leq 4, 1 \leq x_2 \leq 8\}$ for any fixed integer $c \geq 1$. So Y_3^c is the c'th 2-level of Y_3 (ordered by altitude, or third coordinate), and $Y_3^{(j)} = \cup_{t=1}^{j} Y_3^t$ is the 3-dimensional subgrid of Y_2 induced by the first j many 2-levels of Y_3. Further $S_3^t \cong [4 \times 8 \times 4]$ is the 3-dimensional subgrid of Y_3 induced by the set of 2-levels $\{Y_3^c : 4(t-1)+1 \leq c \leq 4t\}$. So $S_3^{(j)} = \cup_{t=1}^{j} S_3^t$ is the 3-dimensional subgrid $[4 \times 8 \times 4j]$ of Y_3 induced by the first $4j$ many 2-levels of Y_3.

We let $u_i(G) = \lceil \frac{|G|}{|\langle Y_{i-1} \rangle|} \rceil = \lceil \frac{|G|}{2^{e_{i-1}}} \rceil$, which we abbreviate with just u_i when G is understood by context or is an arbitrary grid. Since each $(i-1)$-level of Y_i has size $|\langle Y_{i-1} \rangle| = 2^{e_{i-1}}$, u_i is the minimum number of $(i-1)$-levels of Y_i whose union could contain the image $f_i(G)$. Our maps f_i will satisfy $f_i(G) \subseteq Y_i^{(u_i)}$ for each $2 \leq i \leq k$.

We will need the analogue of a t-level for $G = [a_1 \times a_2 \times \cdots \times a_k]$. For $2 \leq i \leq k-1$, an *i-page* of G is any i-dimensional subgrid of G induced by all vertices of G having the same

last $k-i$ coordinate values. Let $P_i = a_{i+1}a_{i+2}\cdots a_k$, which is the number distinct i-pages in G. We define a linear ordering $\prec_i$ on these i-pages as follows. Let D_i and D_i' be two i-pages, with fixed last $k-i$ coordinate values $c_{i+1}, c_{i+2}, \ldots, c_k$ and $c'_{i+1}, c'_{i+2}, \ldots, c'_k$ respectively. Then $D_i \prec_i D_i'$ in this ordering if at the maximum index r, $i+1 \leq r \leq k$, where $c_r \neq c'_r$ we have $c_r < c'_r$. Now index the i-pages of G relative to this ordering by D_i^j, $1 \leq j \leq P_i$, where $r < s$ if and only if $D_i^r \prec_i D_i^s$. Also let $D_i^{(r)} = \bigcup_{j=1}^{r} D_i^j$.

As an example, consider the 4-dimensional grid $H = [3 \times 7 \times 3 \times 2]$, containing 6 many 2-pages each isomorphic to $[3 \times 7]$. For fixed i and j, $1 \leq i \leq 3$ and $1 \leq j \leq 2$, denote by $D_2\langle i,j\rangle$ the 2-page of G given by $D_2\langle i,j\rangle = \{(x_1,x_2,i,j) \in H : 1 \leq x_1 \leq 3, 1 \leq x_2 \leq 7\}$. Then the above ordering of 2-pages of H is given by $D_2^1 = D_2\langle 1,1\rangle \prec_2 D_2^2 = D_2\langle 2,1\rangle \prec_2 D_2^3 = D_2\langle 3,1\rangle \prec_2 D_2^4 = D_2\langle 1,2\rangle \prec_2 D_2^5 = D_2\langle 2,2\rangle \prec_2 D_2^6 = D_2\langle 3,2\rangle$. There are two 3-pages in H, given by $D_3^1 = \{D_2^1, D_2^2, D_2^3\}$ and $D_3^2 = \{D_2^4, D_2^5, D_2^6\}$, and we have $D_3^1 \prec_3 D_3^2$. As this example illustrates, for $3 \leq i \leq k$ the ordering $\prec_{i-1}$ is a refinement of the ordering $\prec_i$ in that if $D_{i-1}^p \subseteq D_i^{p'}$ and $D_{i-1}^q \subseteq D_i^{q'}$ with $p' < q'$, then $p < q$.

From here on we fix $G = [a_1 \times a_2 \times \cdots \times a_k]$ to be a k-dimensional grid. We summarize the above notation, together with selected notation items to be introduced later, in Appendix 2 for convenient reference.

2 The 2-dimensional mapping

Recall the 2-dimensional grid $Y_2 = P(2^{e_1}) \times P(l)$, with $e_1 = \lceil log_2(a_1) \rceil$ and l sufficiently large. In this section we construct a map $f_2 : G \to Y_2^{(u_2)}$, and we abbreviate $m = u_2$ throughout this section, so $m = \lceil \frac{|G|}{2^{e_1}} \rceil$. So f_2 will satisfy $f_2(G) \subseteq P(2^{e_1}) \times P(2^{\lceil log_2(|G|)\rceil - e_1}) \subset Opt(G)$. Additional work will show that for any edge $vw \in E(G)$ we have that $|f_2(v)_1 - f_2(w)_1| + |f_2(v)_2 - f_2(w)_2|$ is small. This map f_2, resembling a map constructed in [9] and [26], will be the first step in an inductive construction leading to a low dilation embedding $f_k : G \to Opt'(G) \subseteq Opt(G)$.

We use the following notation. Let $G(r)$ denote the infinite 2-dimensional grid having r rows, so the vertex and edge sets of $G(r)$ are $V(G(r)) = \{(x,y) \in \mathbb{Z}^2 : 1 \leq x \leq r, 1 \leq y < \infty\}$, and $E(G(r)) = \{(x_1,y_1)(x_2,y_2) : |x_1 - x_2| + |y_1 - y_2| = 1\}$. We let C_i denote the set of vertices (x,y) of $G(r)$ with $x = i$, and refer to this set as "chain i", or the "i'th chain" of $G(r)$.

We can view $V(G)$ as a subset of $V(G(a_1))$ by a natural correspondence $\kappa : V(G) \to V(G(a_1))$ defined as follows. Let $W_i = \prod_{t=2}^{i} a_t$ for $2 \leq i \leq k$. For any vertex $x = (x_1, x_2, \ldots, x_k)$ of G, let $\kappa(x) = (x_1, y)$, where $y = (x_k - 1)W_{k-1} + (x_{k-1} - 1)W_{k-2} + \cdots + (x_3 - 1)W_2 + x_2$. To see the action of κ, let ρ_t be the subset of $V(G)$ consisting of vertices $x \in V(G)$ with $x_1 = t$, $1 \leq t \leq a_1$. Then κ maps the points of ρ_t to the first W_k points of C_t in lexicographic order; that is, if $z = (t, x_2, x_3, \ldots, x_k)$ and $z' = (t, x'_2, x'_3, \ldots, x'_k)$ are two points of ρ_t, then $\kappa(z)_1 = \kappa(z')_1 = t$ and $\kappa(z)_2 < \kappa(z')_2$ if and only if at the largest index r where $x_r \neq x'_r$ we have $x_r < x'_r$. Recall now the ordering D_i^j, $1 \leq j \leq P_i$, of i-pages defined at the end of the last section. Then for fixed $2 \leq i \leq k$ and $1 \leq j \leq P_i$, $\kappa(D_i^j)$ is the subset of points in $G(a_1)$ given by $\{(t,y) : 1 \leq t \leq a_1, (j-1)a_2a_3a_4\cdots a_i + 1 \leq y \leq ja_2a_3a_4\cdots a_i\}$.

Our method in this section is to first construct a low dilation map $f : G(a_1) \to Y_2$. We then obtain the desired f_2 as the composition $f_2 = f \circ \kappa : G \to Y_2^{(m)}$. For the rest of this section we literally identify any point $x \in V(G)$ with $\kappa(x)$, dropping further references to κ

itself. Thus, once $f : G(a_1) \to Y_2$ is constructed, our map $f_2 : G \to Y_2^{(m)}$ will henceforth be viewed as the restriction of f to $G \subset G(a_1)$ (under the identification $x \leftrightarrow \kappa(x)$).

The map $f : G(5) \to Y_2$ is shown in Figure 1(b), and the map $f_2 : [3 \times 7 \times 4 \times a_4] \to Y_2$, obtained by restriction from $f : G(3) \to Y_2$, is shown in Figure 2.

To begin the description of f, let positive integers $1 \le i \le a_1$ and $j \ge 1$ be given. Then f will map either 1 or 2 points of chain C_i to Y_2^j (which recall is column j of Y_2). We encode this information by defining a $(0,1)$ matrix R having a_1 rows and infinitely many columns indexed by the positive integers, where $R_{ij} = 1$ (resp. $R_{ij} = 0$) means that C_i has 2 (resp. 1) image points in Y_2^j under the map f. In the first case these 2 image points are successive in Y_2^j.

Define the first column of R by

$$R_{i1} = \lfloor (\frac{2^{e_1} - a_1}{a_1})i \rfloor - \lfloor (\frac{2^{e_1} - a_1}{a_1})(i-1) \rfloor, 1 \le i \le a_1.$$

For $j > 1$, let $R_{i,j} = R_{i-1,j-1}$ where the row index is viewed modulo a_1. Thus R is just a circulant matrix whose columns are obtained by successive downward shifts of the first column with wraparound, and is illustrated for $a_1 = 5$ in Figure 1(a). The following Lemma shows that the set of 1's in any set of consecutive entries of some row or column of R depends only on the number of such entries, and is one of two successive integers depending on that number.

Lemma 2.1 *The matrix R has the following properties.*
(a) *The sum of entries in any column of R is $2^{e_1} - a_1$.*
(b) *The sum of any t consecutive entries in any row or column of R is either S_t or $S_t + 1$, where $S_t = \lfloor (\frac{2^{e_1} - a_1}{a_1})t \rfloor$.*

Proof. For (a), it suffices to prove the claim for column 1 of R, since any other column of R is just a circular shift of column 1. This column sum is $\sum_{i=1}^{a_1} (\lfloor (\frac{2^{e_1} - a_1}{a_1})i \rfloor - \lfloor (\frac{2^{e_1} - a_1}{a_1})(i-1) \rfloor)$, which telescopes to $2^{e_1} - a_1$.

Consider (b). From the circulant property of R and the constant column sum property from part (a), it suffices to prove this claim for any sum of t consecutive entries (mod a_1) in column 1, say a telescoping sum of the form

$$\sum_{i=r}^{r+t-1} R_{i1} = \lfloor (\frac{2^{e_1} - a_1}{a_1})(r+t-1) \rfloor - \lfloor (\frac{2^{e_1} - a_1}{a_1})(r-1) \rfloor.$$

Now letting $A = (\frac{2^{e_1} - a_1}{a_1})t$ and $B = (\frac{2^{e_1} - a_1}{a_1})(r-1)$, we see that this sum is $\lfloor A + B \rfloor - \lfloor B \rfloor$. But any such difference is either $\lfloor A \rfloor$ or $\lfloor A \rfloor + 1$, so our sum is S_t or $S_t + 1$ as claimed. ∎

The basic idea in constructing f is to fill out $Y_2^{(m)}$ with images under f column by column of $Y_2^{(m)}$ in increasing column index. For any fixed column of $Y_2^{(m)}$, say the j'th column Y_2^j, we proceed down that column mapping points of $G(a_1)$ in nondecreasing chain index. The set $f(C_i) \cap Y_2^j$ will consist of either 1 point or 2 successive points of Y_2^j, depending on whether $R_{ij} = 0$ or 1 respectively.

The construction of f follows. While reading the construction below, the reader may wish to consult Figure 1 illustrating the matrix R and corresponding map $f : G(5) \to Y_2$. For $1 \le r \le a_1$, we let $C^{(r)} = \bigcup_{i=1}^{r} C_i$.

Construction of the map $f : G(a_1) \to Y_2$

1.The images of f in Y_2^1 are given as follows.

(1a) Initial points of C_1 are mapped to Y_2^1 as follows.
If $R_{11} = 0$, then $f(1,1) = (1,1)$ and $c_1 \leftarrow 1$.
If $R_{11} = 1$, then $f(1,1) = (1,1)$, $f(1,2) = (2,1)$, and $c_1 \leftarrow 2$.

(1b) For $i = 1$ to $a_1 - 1$
begin
Assume inductively that $f(C^{(i)}) \cap Y_2^1 = \{(x,1) : 1 \leq x \leq c_i\}$ for an integer c_i. Then initial points of C_{i+1} are mapped to Y_2^1 as follows.
If $R_{11} = 0$, then $f(i+1,1) = (c_i + 1, 1)$, and $c_{i+1} \leftarrow c_i + 1$.
If $R_{11} = 1$, then $f(i+1,1) = (c_i + 1, 1)$, $f_2(i+1,2) = (c_i + 2, 1)$, and $c_{i+1} \leftarrow c_i + 2$.
If $i = a_1 - 1$, then go to step 2.
end

2. For all $1 \leq t \leq N$ (with N sufficiently large), the images of f in Y_2^t are given inductively as follows.

For $j = 1$ to N
begin

Suppose inductively that every point of $Y_2^{(j)}$ has been defined as an image under f. Suppose also that for each $1 \leq i \leq a_1$, we have $C_i \cap f^{-1}(Y_2^{(j)}) = \{(i,t) : 1 \leq t \leq n_{ij}\}$ for suitable integers n_{ij}, with $n_{0j} = n_{i0} = 0$.

[Comment : So the first n_{ij} points of C_i have been mapped to $Y_2^{(j)}$.]

Now define $f(G(a_1)) \cap Y_2^{j+1}$ as follows.
For $r = 0$ to $a_1 - 1$
begin

Suppose inductively that $f(C^{(r)}) \cap Y_2^{j+1}$ is an initial segment of Y_2^{j+1}, say given by $f(C^{(r)}) \cap Y_2^{j+1} = \{(x, j+1) : 1 \leq x \leq c_{r,j+1}\}$, where $c_{0,j} = 0$ for all $j \geq 1$.

Define $f(C_{r+1}) \cap Y_2^{j+1}$ as follows.

(2a) If $R_{r+1,j+1} = 0$, then $f(r+1, 1+n_{r+1,j}) = (c_{r,j+1} + 1, j+1)$, and $c_{r+1,j+1} \leftarrow c_{r,j+1} + 1$.

(2b) If $R_{r+1,j+1} = 1$, and $j+1$ is even then $f(r+1, 1+n_{r+1,j}) = (c_{r,j+1} + 2, j+1)$, $f(r+1, 2+n_{r+1,j}) = (c_{r,j+1} + 1, j+1)$, and $c_{r+1,j+1} \leftarrow c_{r,j+1} + 2$.

(2c) If $R_{r+1,j+1} = 1$ and $j+1$ is odd, then $f(r+1, 1+n_{r+1,j}) = (c_{r,j+1}+1, j+1)$, $f(r+1, 2+n_{r+1,j}) = (c_{r,j+1} + 2, j+1)$, and $c_{r+1,j+1} \leftarrow c_{r,j+1} + 2$.
end
end

Toward analyzing this construction, recall that $m = \lceil \frac{|G|}{2^{e_1}} \rceil$. With the goal of showing (in the theorem which follows) that $f(G) \subseteq Y_2^{(m)}$, we analyze $f^{-1}(Y_2^{(m)})$. Let $C_i(t) = \{(i,y) : 1 \leq y \leq t\}$ be the set of the first t points of chain C_i. Now by steps 2a and 2b, $f(C_i)$ contributes either one point or two successive points to any column Y_2^j, depending on whether $R_{ij} = 0$ or 1 respectively. So $f(C_i)$ contributes exactly $j + \sum_{t=1}^{j} R_{it}$ points to $Y_2^{(j)}$. Thus letting $N_{ij} = j + \sum_{t=1}^{j} R_{it}$, we have $f^{-1}(Y_2^{(m)}) \cap C_i = C_i(N_{im})$. So let $G(a_1, N_{im})$ be the subgraph

of $G(a_1)$ induced by $\bigcup_{i=1}^{a_1} C_i(N_{im})$. The next theorem gives various properties of f, including $Y_2^{(m-1)} \subset f(G) \subseteq f(G(a_1, N_{im})) = Y_2^{(m)}$.

Finally, we define $f_2 : G \to Y_2^{(m)}$ as the restriction of f to the subgraph G of $G(a_1)$. In Figure 2 we illustrate part of $f_2(G)$ for $G = [3 \times 7 \times 4 \times a_4]$ for some $a_4 > 1$. Each 2-page D_2^i of G is isomorphic to $[3 \times 7]$, and the images $f_2(D_2^i)$, $1 \le i \le 4$, are shown in detail with dividers separating the images $f_2(D_2^i)$ and $f_2(D_2^{i+1})$ of successive 2-pages . Near these dividers, for each chain and each 2-page we have placed a box around the image of the chain's first point in that 2-page. For example, under the letter C are three boxed points, representing the images of the first points of each of the three chains in the third 2-page D_2^3. We have also labeled three points in the figure by their preimages in G. For example $(2, 4, 4, ...)$ indicates the preimage $(x_1, x_2, x_3, \cdots, x_k) \in D_2^4 \subset G$ with $x_1 = 2$, $x_2 = 4$, $x_3 = 4$, and so on. There are four 2-pages of G in each 3-page of G, and Figure 2 includes the image $f_2(D_3^1)$ of the first 3-page, consisting of $\bigcup_{i=1}^{4} f_2(D_2^i)$. The beginnings of $f_2(D_2^5)$, this being the first 2-page of the 3-page D_3^2, and of $f_2(D_3^2) = \bigcup_{i=5}^{8} f_2(D_2^i)$ are also illustrated at the far right in the same figure.

Theorem 2.2 *The map $f : G(a_1) \to Y_2$ constructed above has the following properties. Let $m = \lceil \frac{|G|}{2^{e_1}} \rceil$, $S_t = \lfloor (\frac{2^{e_1} - a_1}{a_1})t \rfloor$ (as in Lemma 2.1b), $N_{ij} = j + \sum_{t=1}^{j} R_{it}$, and let $G(a_1, N_{im})$ be the subgraph of $G(a_1)$ induced by $\bigcup_{i=1}^{a_1} C_i(N_{im})$.*

(a) *For each i and j, $|f(C_i) \cap Y_2^j| = 1$ or 2, depending on whether $R_{ij} = 0$ or 1 respectively. Further, if $R_{ij} = 1$, then $f(C_i) \cap Y_2^j$ consists of two successive points of Y_2^j. Also, f is monotone in the sense that $f(i, j)_2 \le f(i, j+1)_2 \le f(i, j)_2 + 1$ for each i and j.*

(b) *Let $L_r(j) = f(C^{(r)}) \cap Y_2^j$. Then $L_r(j)$ is an initial segment, say $\{(d, j) : 1 \le d \le |L_r(j)|\}$, of Y_2^j, with $|L_r(j)| = r + \sum_{i=1}^{r} R_{ij}$.*

(c) *For $i, h, j \ge 1$ with $h \le j$, let $\pi(i, h \to j)$ be the number of points of C_i mapped to columns h through j of $Y_2^{(m)}$. Then $\pi(i, 1, j) = N_{ij}$. Further, for any $r, s \ge 1$ we have $\pi(i, r \to r + j) = j + 1 + S_{j+1}$ or $j + 2 + S_{j+1}$, and $|\pi(i, r \to r + j) - \pi(i, s \to s + j)| \le 1$.*

(d) *For any $1 \le r_1 < r_2 \le a_1$ we have $|N_{r_1,j} - N_{r_2,j}| \le 1$. Also with $L_r(j)$ as in (b), for any $1 \le j_1 < j_2 \le m$ we have $||L_r(j_1)| - |L_r(j_2)|| \le 1$, and $||L_{r+1}(j_1)| - |L_r(j_2)|| \le 2$.*

(e) *$f(G(a_1, N_{im})) = Y_2^{(m)}$.*

(f) *$G \subseteq G(a_1, N_{im})$, and $Y_2^{(m-1)} \subset f(G)$.*

(g) *For any i, j, and r we have $|f(i, r)_2 - f(j, r)_2| \le 1$.*

(h) *For any i, j, and r we have $|f(i, j)_1 - f(i, r)_1| \le 2$.*

(i) *Suppose that $|f(C_r) \cap Y_2^j| = 2$. Then $|L_r(j)| \ge |L_r(j+1)|$ and $N_{r,j} \ge N_{r+1,j}$.*

We omit the involved but straightforward proof of this theorem here, and give it in Appendix 1. The properties listed above can be easily verified in the examples illustrated in Figures 1 and 2.

The following corollary will be used later in proving our dilation bound and the containment $f_2(G) \subseteq Opt(G)$. Its proof also appears in Appendix 1.

Corollary 2.3 *Let v and w be adjacent points of G. The map $f_2 : G \to Y_2^{(m)}$ has the following properties.*

(a) $|f_2(v)_1 - f_2(w)_1| \leq 3$, *and* $|f_2(v)_2 - f_2(w)_2| \leq 1$.

(b) $|f_2(v)_1 - f_2(w)_1| + |f_2(v)_2 - f_2(w)_2| \leq 3$.

(c) $f_2(G) \subseteq Opt(G)$.

(d) *Let* T *and* T' *be segments of* p *consecutive points on chains* C_i *and* C_j *respectively,* $1 \leq i, j \leq a_1$, *where possibly* $i = j$. *Let* c *and* c' *be the number of columns of* Y_2 *spanned by* $f_2(T)$ *and* $f_2(T')$ *respectively. Then* $|c - c'| \leq 1$.

(e) *For* $1 \leq r \leq P_2$, *let* $r' = min\{c : f_2(D_2^{(r)}) \subseteq Y_2^{(c)}\}$. *Then* $Y_2^{(r'-1)} \subset f_2(D_2^{(r)})$ *and* $|Y_2^{(r')} - f_2(D_2^{(r)})| < 2^{e_1}$.

3 The idea of the general construction, with examples

In this section we give the idea behind our general construction, saving complete details and proofs of validity for later sections. We continue with the notation of Section 1.2.

The overall plan is to construct a sequence of maps $f_i : G \to [\langle Y_{i-1}\rangle \times P(u_i)] = Y_i^{(u_i)}$, $u_i = \lceil \frac{|G|}{|\langle Y_{i-1}\rangle|} \rceil = \lceil \frac{|G|}{2^{e_{i-1}}} \rceil$, $2 \leq i \leq k$, the first of which is f_2 from section 2. Since $u_i \leq 2^{\lceil log_2(|G|)\rceil - e_{i-1}}$, we will have $f_i(G) \subseteq Y_i^{(u_i)} \subseteq \langle Y_{i-1}\rangle \times P(2^{\lceil log_2(|G|)\rceil - e_{i-1}})$. The last graph is a spanning subgraph of $Opt(G)$, so $f_i(G) \subseteq Opt(G)$ for $2 \leq i \leq k$. In particular, for $i = k$ we get $f_k(G) \subseteq Opt'(G)$. The maps f_i will be successive approximations to f_k in that for any $x \in V(G)$ and $2 \leq i < k$ we will have $f_i(x)_{1\to i-1} = f_k(x)_{1\to i-1}$.

The final map f_k gives the basic geometry of our construction. We then apply a labeling L of the points of $Opt'(G)$ with hypercube addresses from $Opt(G)$ to obtain the final embedding $H^k : G \to Opt(G)$, where $H^k = L \circ f_k$. We construct the maps f_i inductively, letting f_{i+1} be the composition $f_{i+1} = \sigma_i \circ I_i \circ f_i$, using maps I_i and σ_i described below. Recall that $P_i = a_{i+1}a_{i+2}\cdots a_k$ is the number of i-pages in G.

So suppose $f_i : G \to Y_i^{(u_i)}$ has been constructed, and we outline the construction of $f_{i+1} : G \to Y_{i+1}^{(u_{i+1})}$. Now $I_i : Y_i^{(u_i)} \to S_i^{(P_i)} \subset Y_i$ is a one to one "inflation" map which spreads out the image $f_i(G)$ "evenly" in $S_i^{(P_i)}$ by successively "skipping over" certain carefully chosen $(i-1)$-levels of Y_i that are designated "blank". See Figure 3 for an example where $i = 2$, and where f_2 is the map of Figure 2 and blank 1-levels (columns) are shaded.

We let $S_i^{(P_i)}(G)$ be the set of points in $S_i^{(P_i)}$ lying in nonblank $(i-1)$-levels (i.e. levels not designated "blank") of $S_i^{(P_i)}$. We stipulate that $(I_i \circ f_i)(G) \subset S_i^{(P_i)}(G)$. The number of successive i-sections S_i^r in the range of I_i is the same as the number of i-pages D_i^r in G, each equalling P_i, and for each r we associate S_i^r with D_i^r in a sense to be made clear below.

Define $I_i : Y_i^{(u_i)} \to S_i^{(P_i)}$ as follows, for now assuming that certain $(i-1)$-levels of Y_i (all lying within $S_i^{(P_i)}$) have been designated blank. For $x = (x_1, x_2, \ldots, x_i) \in Y_i^{(u_i)}$, we let $I_i(x) = (x_1, x_2, \ldots, x_{i-1}, x_i')$, where x_i' is the common i-coordinate in the x_i'th nonblank $(i-1)$-level of Y_i (in order of increasing i-coordinate). Now for any $S \subseteq Y_i^{(u_i)}$, let $I_i(S) = \{I_i(s) : s \in S\}$. Observe that by its definition, I_i preserves $(i-1)$-levels; that is $I_i(Y_i^t) = Y_i^{t'}$ is an $(i-1)$-level of Y_i. Also for $1 \leq s < t \leq u_i$, where $I_i(Y_i^s) = Y_i^{s'}$ and $I_i(Y_i^t) = Y_i^{t'}$, we have $s' < t'$; that is, I_i preserves the order (by increasing i-coordinate) of $(i-1)$-levels. We can picture I_i as an order preserving spreading out of the u_i many $(i-1)$-levels of $Y_i^{(u_i)}$ containing $f_i(G)$ among

the $P_i 2^{e_i - e_{i-1}}$ many $(i-1)$-levels of $S_i^{(P_i)}$. The image $I_i(Y_i^{(u_i)})$ becomes the set of nonblank $(i-1)$-levels (u_i of them) in $S_i^{(P_i)}$. The remaining $P_i 2^{e_i - e_{i-1}} - u_i$ many $(i-1)$-levels of $S_i^{(P_i)}$ are blank, and distributed among the nonblank $(i-1)$-levels so that certain balance properties outlined below are satisfied.

Consider the example $G' = [3\times 7\times 4\times 3]$. Start with $f_2 : G' \to Y_2^{(u_2(G'))}$ given in the previous section, where $u_2(G') = \lceil \frac{252}{4} \rceil = 63$. Figure 2 gives the initial part of $f_2(G')$, while Figures 3a) and b), illustrate the map $I_2 : Y_2^{(u_2(G'))} \to S_2^{(P_2)}$, $P_2 = 12$. Each 2-section S_2^j, $1 \le j \le 12$, satisfies $S_2^j \cong \langle Y_1 \rangle \times P(2^{e_2 - e_1}) = P(4) \times P(8)$. So $S_2^{(P_2)}$ has $P_2 2^{e_2 - e_1} = 12 \cdot 8 = 96$ columns (i.e. $(i-1)$-levels where $i = 2$), of which the 63 nonblank ones comprising $I_2(Y_2^{(63)})$ contain $(I_2 \circ f_2)(G')$, while the remaining 33 blank ones (shaded) contain no points of $(I_2 \circ f_2)(G')$. Note in the figure how I_2 preserves the order in $S_2^{(12)}$ of the 63 columns in the domain $Y_2^{(63)}$ of I_2, and the 33 blank columns (shaded) of $S_2^{(12)}$ are distributed throughout $S_2^{(12)}$.

Turning to arbitrary G, the quantity and distribution of blank $(i-1)$-levels within $S_i^{(P_i)}$ will be such that for each $1 \le r \le P_i$, the subgraph $S_i^{(r)}$ of Y_i has barely enough nonblank $(i-1)$-levels to host $(I_i \circ f_i)(D_i^{(r)})$. In effect, we want $S_i^{(r)}$ to have at least as much nonblank volume as $|D_i^{(r)}|$ for each $r \ge 1$, but barely so in increments the size of an $(i-1)$-level in Y_i. We formulate this condition precisely as follows. Let $s_i(j)$ be the number of blank $(i-1)$-levels in section S_i^j. Observing that $|D_i^j| = a_1 a_2 \cdots a_i$ for any j, that each $(i-1)$-level of Y_i has $2^{e_{i-1}}$ points, and that each section S_i^j consists of $2^{e_i - e_{i-1}}$ many pairwise disjoint $(i-1)$-levels, we require that

$$\lceil \frac{r a_1 a_2 \cdots a_i}{2^{e_{i-1}}} \rceil + \sum_{j=1}^{r} s_i(j) = r 2^{e_i - e_{i-1}} \tag{1}$$

for each $1 \le r \le P_i$.

Next, the map $\sigma_i : (I_i \circ f_i)(G) \to \langle Y_i \rangle \times P(u_{i+1}) = Y_{i+1}^{(u_{i+1})}$ "stacks" the sets $S_i^j \cap (I_i \circ f_i)(G)$, $1 \le j \le P_i$, over the single section $S_i^1 \cong \langle Y_i \rangle$ as follows. Let $x = (x_1, x_2, \ldots, x_i) \in S_i^j \cap (I_i \circ f_i)(G)$, $1 \le j \le P_i$. Let $\bar{x}_i \equiv x_i \pmod{2^{e_i - e_{i-1}}}$, $1 \le \bar{x}_i \le 2^{e_i - e_{i-1}}$, be the congruence class of x_i mod $2^{e_i - e_{i-1}}$. Define the first i coordinates of $\sigma_i(x)$ by $\sigma_i(x)_{1\to i} = (x_1, x_2, \ldots, x_{i-1}, \bar{x}_i)$. Observe that since $x \in S_i^j$, we have $x_i = (j-1)2^{e_i - e_{i-1}} + \bar{x}_i$, and that $\sigma_i(x)_{1\to i} \in S_i^1$. To get the $i+1$'st (and last) coordinate, let c be the number of points $y = (y_1, y_2, \ldots, y_i) \in S_i^t \cap (I_i \circ f_i)(G)$, $1 \le t \le j$, satisfying $\sigma_i(y)_{1\to i} = \sigma_i(x)_{1\to i}$. Then define $\sigma_i(x) = (x_1, x_2, \ldots, x_{i-1}, \bar{x}_i, c)$.

Finally define f_{i+1} as the composition $f_{i+1} = \sigma_i \circ I_i \circ f_i$.

For a fixed point $(x_1, x_2, \ldots, x_{i-1}, \bar{x}_i) \in S_i^1$, we view the set of images $(x_1, x_2, \ldots, x_{i-1}, \bar{x}_i, c)$ under σ_i as a stack addressed by $(x_1, x_2, \ldots, x_{i-1}, \bar{x}_i)$ extending into the $(i+1)$'st dimension. Thus each point $(x_1, x_2, \ldots, x_{i-1}, \bar{x}_i)$ of S_i^1 becomes the address of such a stack, and the image $\sigma_i(x) = (x_1, x_2, \ldots, x_{i-1}, \bar{x}_i, c)$ is the c'th point "up" in this stack. See Figure 4, to which we return later with a full explanation, for an initial look at σ_2.

Since the domain of σ_i is $(I_i \circ f_i)(G)$, which is a set contained in the collection of nonblank $(i-1)$-levels of $S_i^{(P_i)}$, it follows that the points in blank $(i-1)$-levels of $S_i^{(P_i)}$ make no contribution under the map σ_i to the aforementioned stacks. We can picture the images $\sigma_i(x)$, $x \in S_i^j \cap (I_i \circ f_i)(G)$, as "falling through" blank $(i-1)$-levels $Y_i^d \subset S_i^t, t < j$, with $d \equiv x_i \pmod{2^{e_i - e_{i-1}}}$. But if such a Y_i^d is nonblank, and if $z = (x_1, x_2, \cdots, x_{i-1}, d) \in Y_i^d \cap (I_i \circ f_i)(G)$, then $\sigma_i(z)_{1\to i} = \sigma_i(x)_{1\to i}$. Thus z does contribute (under σ_i) to the same stack as x; that

is, both $\sigma_i(z)$ and $\sigma_i(x)$ belong to the stack addressed by $(x_1, x_2, \ldots, x_{i-1}, \bar{x}_i) \in S_i^1$. Further, $\sigma_i(z)_{i+1} < \sigma_i(x)_{i+1}$ by definition of c and since $t < j$; that is, $\sigma_i(z)$ appears "below" $\sigma_i(x)$ in this stack. So we see that each section S_i^j, $1 \le j < P_i$, contributes (via σ_i) either 0 or 1 point to the stack $(x_1, x_2, \ldots, x_{i-1}, \bar{x}_i) \in S_i^1$. It contributes 1 point precisely when its unique $(i-1)$-level $Y_i^d \subset S_i^j$ satisfying $d \equiv \bar{x}_i \pmod{2^{e_i - e_{i-1}}}$ is not blank and the point $z = (x_1, x_2, \cdots, x_{i-1}, d) \in Y_i^d$ of S_i^j lies in $(I_i \circ f_i)(G)$.

As an example, consider $G'' = [3 \times 7 \times 4] \subset G'$. We construct $f_3(G'') = (\sigma_2 \circ I_2 \circ f_2)(G'')$. Note that $e_1 = 2$, $e_2 = 5$, $P_2 = 4$, $u_2(G'') = \lceil \frac{84}{4} \rceil = 21$, and $u_3(G'') = \lceil \frac{84}{32} \rceil = 3$. Again each 2-section S_2^j, $1 \le j \le P_2 = 4$, is isomorphic to $P(2^{e_1}) \times P(2^{e_2 - e_1}) = P(4) \times P(8)$, and $S_2^{(P_2)}$ has $P_2 \cdot 2^{e_2 - e_1} = 4 \cdot 8 = 32$ columns.

Step 1: Start with $f_2 : G'' \to Y_2^{(u_2(G''))} = Y_2^{(21)}$ given in section 2 and illustrated as $f_2(D_3^1)$ in Figure 2.

Step 2: Perform the map $I_2 : Y_2^{(21)} \to S_2^{(P_2)} = S_2^{(4)}$ by letting $I_2(a, b) = (a, b')$, where b' is the b'th nonblank column of $S_2^{(P_2)}$, ordered by increasing second coordinate. The result is shown in Figure 3a), where $I_2(f_2(G''))$ is unshaded and blank columns are shaded. The choice of blank columns will be discussed later.

On comparing with Figure 2, we see that $I_2(f_2(G''))$ is indeed obtained from $f_2(G'')$ by skipping over the designated blank columns. Each 2-section $S_2^i \subset S_2^{(4)}$, $1 \le i \le 4$, is the host for the 2-page D_2^i of G'', with the possible exception of the images of first points of chains in D_2^i (these being boxed in Figure 3) which may appear in the last nonblank column of S_2^{i-1} (instead of in S_2^i).

Step 3: Perform the stacking map $\sigma_2 : (I_2 \circ f_2)(G'') \to Y_3^{(u_3(G''))} = Y_3^{(3)}$. Thus for each $(a, b) \in S_2^j \cap (I_2 \circ f_2)(G'')$, we let $\sigma_2((a, b)) = (a, \bar{b}, c)$, where $\bar{b} \equiv y \pmod 8$, $1 \le \bar{b} \le 8$, and $c = |\{(a, b') \in S_2^t \cap (I_2 \circ f_2)(G'') : t \le j,\ b' \equiv b \pmod 8\}|$. As above, we view $(a, \bar{b})$ as the address in S_2^1 of the stack on which (a, b) has been placed by σ_2, and $\sigma_2((a, b))$ is the c'th point "up" on this stack.

In Figure 4 we illustrate $f_3(G'') = \sigma_2((I_2 \circ f_2)(G''))$; that is, how σ_2 stacks the four sets $S_2^j \cap (I_2 \circ f_2)(G'')$, $1 \le j \le 4$, over $S_2^1 = \langle Y_2 \rangle$ to yield $f_3(G'')$. At top center we begin with $I_2(Y_2^{(21)}) \subseteq S_2^{(P_2)} = S_2^{(4)}$ where the four 2-sections S_2^j, $1 \le j \le 4$ comprising $S_2^{(4)}$, are placed vertically in succession for convenience. We now perform the stacking, with the result shown at lower left in the figure. Here we regard the bottom 4×8 layer (of the three layers in the result) as a copy of S_2^1, each of whose 32 points is the address of a stack. Consider the 4 points lying in column 2 of this S_2^1. As shown in the figure, each of these points is the address of a stack of height 3. For brevity let us write $\sigma_2(S_2^j)$ for $\sigma_2(S_2^j \cap (I_2 \circ f_2)(G''))$. The contributions to any one of these 4 stacks come from $\sigma_2(S_2^1)$, $\sigma_2(S_2^3)$, and $\sigma_2(S_2^4)$ (indicated respectively by the labels $1, 3, 4$ in these stacks). To see how this happens, refer back to the top center of this figure and look at the points in column 2 in the 4 sections S_2^j, $1 \le j \le 4$, being stacked. Since column 2 of the second of these, S_2^2, is blank, the points in it make no contribution to the stacks whose addresses lie in column 2 of S_2^1 at lower left. But the points in column 2 of S_2^1, S_2^3, and S_2^4 (at top center) lie in nonblank columns and are contained in $(I_2 \circ f_2)(G'')$. Hence they do contribute to the column 2 stacks of S_2^1 at lower left. We can view such points in column 2 of S_2^3 and S_2^4 in the top center as "falling through" blank column 2 in S_2^2 under the action of σ_2. Similarly every point in column 1 of $S_2^1 = \langle Y_2 \rangle$ at lower left is the address of a stack of height 2. The contributions to these stacks are from $\sigma_2(S_2^2)$, $\sigma_2(S_2^3)$, indicated

by labels 2 and 3 in these stacks. These contributions "fall through" blank column 1 of S_2^1 (at top center) under the action of σ_2. The lower right of the figure shows how individual image points are affected by this stacking. For example, the images under σ_2 of paths in $S_2^2 \cap (I_2 \circ f_2)(G'')$ (in bold at top center) jump between levels of the final result at lower right. Note also that the maximum stack height is indeed $u_3(G'') = 3$, achieved at stacks addressed by points in columns $2, 3, 5, 6$, and 8 of S_2^1. So $f_3(G'') \subseteq Y_3^{(3)}$.

In the rest of this section we show how to assign blank columns in $S_2^{(P_2)}$ (used in constructing I_2 and then f_3), and generally how to assign blank $(i-1)$-levels in $S_i^{(P_i)}$ (used in constructing I_i and then f_{i+1}) for an arbitrary multidimensional grid G. We will see that the ability to make this assignment is implied by the existence of a certain class of $(0,1)$ matrices, whose construction we give in the next section. For now we describe conditions which our assignment will satisfy that are sufficient for making our embedding f_k (and later H^k) have the required dilation and containment properties. To start, consider the assignment of blank columns in $S_2^{(P_2)}$.

(a) The number $s_2(j)$ of blank columns in each 2-section S_2^j, $1 \le j \le P_2$, is chosen so that after $(I_2 \circ f_2)(G)$ "skips over" these columns, each subgraph $S_2^{(r)}$ (the union of the first r many 2-sections), $1 \le r \le P_2$, has barely enough nonblank columns to host the image $(I_2 \circ f_2)(D_2^{(r)})$ of the first r many 2-pages of G. This is expressed in equation (1) in the case $i = 2$, which says that $\lceil \frac{ra_1a_2}{2^{e_1}} \rceil + \sum_{j=1}^{r} s_2(j) = r2^{e_2-e_1}$ for each $1 \le r \le P_2$. Thus each image $(I_2 \circ f_2)(D_2^j)$ lies in its own section S_2^j, apart from images of first points of chains in D_2^j as explained previously and shown in Figure 3. In the example $G'' = [3 \times 7 \times 4]$, the sequence $s_2(j)$ can be found recursively from equation (1) (with $i = 2$) to be $s_2(1) = 2$, $s_2(2) = 3$, $s_2(3) = 3$, and $s_2(4) = 3$, and these numbers of blank columns are shown in the four sections respectively in Figure 3a).

(b) The blank columns are distributed over the various 2-sections S_2^j so that for $xy \in E(G)$ the contribution to $dist_{Y_k}(f_k(x), f_k(y))$ from $|f_k(x)_2 - f_k(y)_2|$ is small. As background, note that $dist_{Y_k}(f_k(x), f_k(y)) = \sum_{i=1}^{k} |f_k(x)_i - f_k(x)_i|$. It turns out that f_k will satisfy $|f_k(x)_i - f_k(x)_i| = |f_{i+1}(x)_i - f_{i+1}(x)_i| = |(I_i \circ f_i)(x)_i - (I_i \circ f_i)(y)_i|$ for each $1 \le i \le k-1$, where the last difference is taken mod $2^{e_i - e_{i-1}}$. The last equality for $i = 2$ can be verified in the construction of $f_3(G'')$ and figures above.

Focusing on the case $i = 2$, suppose that x and y agree in their first two coordinates. So x and y are corresponding points in their respective 2-pages, say $x \in D_2^s$ and $x \in D_2^t$. Now we wish to keep $|(I_2 \circ f_2)(x)_2 - (I_2 \circ f_2)(y)_2|$ small mod $2^{e_2-e_1}$. Thus we want $(I_2 \circ f_2)(x)$ to be skipping over roughly the same number of blank columns in S_2^s as does $(I_2 \circ f_2)(y)$ in S_2^t. To accomplish this, given that s and t are arbitrary as are x and y as corresponding points, we will require a strong balance, over all 2-sections S_2^j, in the frequency of blank columns in any initial segment of columns in S_2^j.

A similar balance will be required in the frequency of blank $(i-1)$-levels in any initial segment of $(i-1)$-levels of any i-section S_i^j. A precise formulation of this requirement with additional detail will be described below.

(c) The assignment of blank columns across all 2-sections S_2^j, $1 \le j \le P_2$, is uniformly distributed mod $2^{e_2-e_1}$. This condition allows us to stack the sections S_2^j on top of each other, over a single 2-section $S_2^1 = \langle Y_2 \rangle$, so that the stacks addressed in S_2^1 have roughly equal stack

heights. In the example $G'' = [3 \times 7 \times 4]$ illustrated in Figure 3a, note that the 32 columns in 2-sections S_2^j, $1 \le j \le 4$, can be partitioned into congruence classes mod 8 by column number, and each congruence class has exactly 1 or 2 of its columns designated blank. The result is that stack heights differing by at most 1.

The goals described above in (a)-(c) can be formulated as combinatorial conditions to be satisfied by the designation of blank columns, and generally of blank $(i-1)$-levels. Starting with the designation of blank columns, it will be convenient to define a $P_2 \times 2^{e_2-e_1}$, $(0,1)$ matrix $F(2) = (f_{cd}(2))$. The rows of $F(2)$ correspond to the 2-sections S_2^c, $1 \le c \le P_2$, and the columns of $F(2)$ to the columns ($2^{e_2-e_1}$ of them) within each 2-section. We let $f_{cd}(2) = 1$ if column d in section S_2^c (which recall is column $Y_2^{(c-1)(2^{e_2-e_1})+d}$ in Y_2) is blank, and $f_{cd}(2) = 0$ if that column is nonblank. Since $s_2(c)$ is the number of blank columns in S_2^c, the sum of entries in row c of $F(2)$ is

$$\sum_{d=1}^{2^{e_2-e_1}} f_{cd}(2) = s_2(c). \tag{2}$$

for $1 \le c \le P_2$.

Toward formulating the goal expressed in (c), consider now the contribution, through the stacking map σ_2, from $S_2^{(r)}$ to any stack addressed by a point in S_2^1. For any stack address $(x,y) \in S_2^1$, let $Stack_3((x,y),r)) = \{\sigma_2(z) : \sigma_2(z)_{1\to 2} = (x,y), z \in S_2^{(r)}\}$, which is the set of points in the stack addressed by (x,y) whose preimages under the map σ_2 come from $S_2^{(r)}$. Now for any $\sigma_2(z) \in Stack_3((x,y),r))$, we have $z = (x,d)$, with $d \equiv y \pmod{2^{e_2-e_1}}$, and column Y_2^d is nonblank. Then for $r < P_2$ we see that the stack height $|Stack_3((x,y),r))|$ is the number of zeros of the matrix $F(2)$ lying in column y and within rows 1 through r . So to keep stack heights nearly equal over all stack addresses $(x,y) \in S_2^1$, we require that this number of zeros is nearly the same over all columns y in $F(2)$. For this, it suffices to have the number of 1's in rows 1 through r of any column nearly the same; that is, to have nearly equal initial column sums. This becomes the condition

$$|\sum_{c=1}^{r} f_{cy}(2) - \sum_{c=1}^{r} f_{cy'}(2)| \le 1. \tag{3}$$

for any $1 \le y, y' \le 2^{e_2-e_1}$ and $1 \le r \le P_2$. It says that the blank columns are uniformly distributed mod $2^{e_2-e_1}$.

To formulate (b), for integers a and b let $||a-b||$ be the difference $a-b$ taken mod $2^{e_2-e_1}$. Let $xy \in E(G)$, say with $x \in D_2^s$ and $y \in D_2^t$, $s \ne t$. Since x and y agree in their first two coordinates, x and y are each the p'th points in D_2^s and D_2^t of their respective chains, for some $1 \le p \le a_2$. By equation (1) for $i = 2$, $S_2^{(r)}$ has barely enough nonblank columns to contain $(I_2 \circ f_2)(D_2^{(r)})$ for any $1 \le r \le P_i$. Thus by the monotonicity property of f_2 in Theorem 2.2a, for any $j \ge 1$, the image under $I_2 \circ f_2$ of the first point of any chain in D_2^j must lie either in the last nonblank column of S_2^{j-1} or the first nonblank column of S_2^j. Now let T (resp. (T')) be the set of the first p points in D_2^s (resp. D_2^t) in the chain containing x (resp. y). Also let c (resp. c') be the number of columns of Y_2 spanned by $f_2(T)$ (resp. $f_2(T')$), and thus the number of nonblank columns of Y_2 spanned by $(I_2 \circ f_2)(T)$ (resp. $(I_2 \circ f_2)(T')$). By Corollary 2.3d we have $|c - c'| \le 1$. Thus the contribution to $||(I_2 \circ f_2)(x)_2 - (I_2 \circ f_2)(y)_2||$

from the difference between the number of nonblank columns in S_2^s (resp. S_2^t) preceding $(I_2 \circ f_2)(x)$ (resp. $(I_2 \circ f_2)(y)$) is small (in fact ≤ 1). The same contribution due to the difference in starting columns of $(I_2 \circ f_2)(T)$ and $(I_2 \circ f_2)(T')$ is also small (≤ 1) by the above. Thus $||(I_2 \circ f_2)(x)_2 - (I_2 \circ f_2)(y)_2||$ depends primarily on the number N_1 (resp. N_2) of blank columns in S_2^s (resp. S_2^t) preceding the column containing $(I_2 \circ f_2)(x)$ (resp. $(I_2 \circ f_2)(y)$). Each column counted by N_1 (resp. N_2) pushes the image $(I_2 \circ f_2)(x)$ (resp. $(I_2 \circ f_2)(y)$) one more column to the right in S_2^s (resp. S_2^t). So we want to keep $|N_1 - N_2|$ small. Since each blank column corresponds to a 1 in $F(2)$, we see that each of N_1 and N_2 is just an initial row sum in $F(2)$ (row s for N_1 and row t for N_2). These considerations motivate the goal of keeping the difference between corresponding initial row sums in $F(2)$ small. It will suffice for our purposes to have

$$|\sum_{j=1}^{b} f_{sj}(2) - \sum_{j=1}^{b} f_{tj}(2)| \leq 2. \tag{4}$$

for any, $1 \leq s, t \leq a_3 a_4 \cdots a_k$, $1 \leq b \leq 2^{e_2 - e_1}$.

For fixed i, we will see that the sequence $\{s_i(j)\}$, $1 \leq j \leq P_i$, recursively defined by (1) satisfies $|s_i(j_1) - s_i(j_2)| \leq 1$ for all $1 \leq j_1, j_2 \leq P_i$. Applying this to the case $i = 2$, we can view the satisfying of conditions (a)-(c), as relying on the construction of a $P_2 \times 2^{e_2 - e_1}$, $(0,1)$ matrix $F(2)$ with prescribed row sums $s_2(c)$, $1 \leq c \leq P_i$, these sums differing by at most 1 (from the preceding sentence and (2)). Further, $F(2)$ will have balanced initial column sums and balanced initial row sums (from (3) and (4)).

Such an $F(2)$ for the embedding $f_3 : G'' = [3 \times 7 \times 4] \to Y_3^{(3)}$ discussed above, where $F(2)$ has $P_2 = 4$ rows, is illustrated by the $(0,1)$ matrix in the right of Table 1a). The fractional matrices at left from which this and the other $(0,1)$ matrices in this table are derived will be explained later. This $F(2)$ encodes which columns are designated 'blank' in performing the inflation step $I_2 : Y_2^{(u_2(G''))} \to S_2^{(4)}$. The stacking map σ_2 is then applied to yield the final embedding $f_3 = \sigma_2 \circ I_2 \circ f_2 : [3 \times 7 \times 4] \to Y_3^{(3)}$.

The corresponding requirements for arbitrary dimension $i \geq 2$ are analogous. We return to the construction of f_{i+1} from f_i as the composition $f_{i+1} = \sigma_i \circ I_i \circ f_i$. Recall that there are $P_i = a_{i+1} a_{i+2} \ldots a_k$ many i-sections S_i^j, $1 \leq j \leq P_i$, and each such i-section has $2^{e_i - e_{i-1}}$ many $(i-1)$-levels. So let $F(i) = (f_{cd}(i))$ be the $P_i \times 2^{e_i - e_{i-1}}$, $(0,1)$ matrix (analogous to $F(2)$) defined by $f_{cd}(i) = 1$ if the d'th $(i-1)$-level of i-section S_i^c (that is, $(i-1)$-level $Y_i^{(c-1)2^{e_i - e_{i-1}} + d}$ of Y_i) is blank, and $f_{cd}(i) = 0$ otherwise. So we require the analogues of the relations (2 - 4);

$$\sum_{j=1}^{2^{e_i - e_{i-1}}} f_{cj}(i) = s_i(c), \tag{5}$$

for $1 \leq c \leq P_i$,

$$|\sum_{c=1}^{r} f_{cy}(i) - \sum_{c=1}^{r} f_{cy'}(i)| \leq 1, \tag{6}$$

for $1 \leq y, y' \leq 2^{e_i - e_{i-1}}$, $1 \leq r \leq P_i$,

$$|\sum_{d=1}^{r} f_{sd}(i) - \sum_{d=1}^{r} f_{td}(i)| \leq 2, \tag{7}$$

for $1 \leq s, t \leq P_i$, $1 \leq r \leq 2^{e_i - e_{i-1}}$.

Relation (5) says that there are $s_i(c)$ many blank $(i-1)$-levels in S_i^c. Relation (6) gives balanced initial column sums in $F(i)$. This implies that blank $(i-1)$-levels are uniformly distributed mod $2^{e_i - e_{i-1}}$. This ensures balanced stack heights, under the map σ_i, for stacks addressed by $S_i^1 \cong \langle Y_i \rangle$. Finally (7) will imply (after some work) that $||(I_i \circ f_i)(x)_i - (I_i \circ f_i)(y)_i||$ is small for any $xy \in E(G)$, with $x \in D_i^s$ and $x \in D_i^t$, $1 \leq s, t \leq P_i$. This will make the contribution to $dist_{Y_k}(f_k(x), f_k(x))$ from $|f_k(x)_i - f_k(y)_i|$ small for corresponding points (x and y) in distinct i-pages. Applying this requirement for all i will keep $dilation(f_k)$ small.

We are thus reduced to the construction of a $(0,1)$ matrix $F(i)$ for each $2 \leq i \leq k-1$, with prescribed row sums $s_i(j)$ (with values recursively computed using (1)), $1 \leq j \leq P_i$, differing by at most 1, and balanced initial column and row sums as required in (6) and (7). We construct such matrices in the next section.

Returning to $G' = [3 \times 7 \times 4 \times 3]$ we illustrate the construction of $f_4 : G' \to Y_4^{(u_4(G'))} = Y_4^{(2)}$. We have $u_2(G') = \lceil \frac{|G'|}{2^{e_1}} \rceil = \lceil \frac{252}{4} \rceil = 63$, $u_3(G') = \lceil \frac{|G'|}{2^{e_2}} \rceil = \lceil \frac{252}{32} \rceil = 8$, and $u_4(G') = \lceil \frac{|G'|}{2^{e_3}} \rceil = \lceil \frac{252}{128} \rceil = 2$.

The maps f_2, f_3, and f_4 are constructed in succession. The map $f_2 : G' \to Y_2^{(63)}$ is given in the previous section. To build $f_3 = \sigma_2 \circ I_2 \circ f_2$ the next step is to define the inflation map I_2. For this, note that $P_2 = 12$, so there will be 12 many 2-sections S_2^j, $1 \leq j \leq 12$, each isomorphic to $P(4) \times P(8)$. Thus I_2 has the form $I_2 : Y_2^{(63)} \to S_2^{(12)}$. The sequence $\{s_2(j)\}$, $1 \leq j \leq 12$, is calculated inductively using (1) for $i = 2$, with $e_1 = 2$ and $e_2 = 5$, to give $2, 3, 3, 3, 2, 3, 3, 3, 2, 3, 3, 3$. A balanced distribution of blank columns among the sections S_2^j, $1 \leq j \leq 12$ satisfying (1) - (4) is given by the 12×8, $(0,1)$ matrix $F(2)$ shown at right in Table 1b). So I_2 will make make $f_2(G')$ skip over the blank columns in $S_2^{(12)}$ as in the previous example. The blank columns (shaded) among the first 4 sections S_2^1, S_2^2, S_2^3, S_2^4, as encoded by the first 4 rows of the matrix in the right of Table 1b), are illustrated in Figure 3a. The blank columns among the remaining 8 sections S_2^j, $5 \leq j \leq 12$, (encoded by the last 8 rows of the same matrix) are illustrated in Figure 3b), where these 8 sections are surrounded by a box. We put small boxes around images of first points of chains within each 2-page.

$$\begin{bmatrix} \mathbf{1/4} & \mathbf{1/4} & \dots & \mathbf{1/4} \\ \mathbf{3/8} & \mathbf{3/8} & \dots & \mathbf{3/8} \\ \mathbf{3/8} & \mathbf{3/8} & \dots & \mathbf{3/8} \\ \mathbf{3/8} & \mathbf{3/8} & \dots & \mathbf{3/8} \end{bmatrix} \quad \begin{matrix} row \\ sum \\ 2 \\ 3 \\ 3 \\ 3 \end{matrix} \quad \begin{matrix} Knuth-like \\ rounding \\ \to \end{matrix} \quad \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \end{bmatrix}$$

a) Matrix $F(2)$ encoding blank columns for the embedding $f_3 : [3 \times 7 \times 4] \to Y_3$

$$\left[\begin{array}{cccc} \mathbf{1/4} & \mathbf{1/4} & \ldots & \mathbf{1/4} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \\ \hline \mathbf{1/4} & \mathbf{1/4} & \ldots & \mathbf{1/4} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \\ \hline \mathbf{1/4} & \mathbf{1/4} & \ldots & \mathbf{1/4} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \\ \mathbf{3/8} & \mathbf{3/8} & \ldots & \mathbf{3/8} \end{array}\right] \begin{array}{c} 2 \\ 3 \\ 3 \\ 3 \\ 2 \\ 3 \\ 3 \\ 3 \\ 2 \\ 3 \\ 3 \\ 3 \end{array} \quad \begin{array}{c} Knuth-like \\ rounding \\ \rightarrow \end{array} \quad \left[\begin{array}{cccccccc} 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ \hline 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ \hline 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 \end{array}\right]$$

b) Matrix $F(2)$ encoding blank columns for the embedding $f_3 : [3 \times 7 \times 4 \times 3] \to Y_3$

$$\left[\begin{array}{cccc} \mathbf{1/4} & \mathbf{1/4} & \mathbf{1/4} & \mathbf{1/4} \\ \mathbf{1/4} & \mathbf{1/4} & \mathbf{1/4} & \mathbf{1/4} \\ \mathbf{1/2} & \mathbf{1/2} & \mathbf{1/2} & \mathbf{1/2} \end{array}\right] \begin{array}{c} 1 \\ 1 \\ 2 \end{array} \quad \begin{array}{c} Knuth-like \\ rounding \\ \rightarrow \end{array} \quad \left[\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{array}\right]$$

c) Matrix $F(3)$ encoding blank 2-levels for the embedding $f_4 : [3 \times 7 \times 4 \times 3] \to Y_4$

Table 1: Matrices encoding blank levels

Next we apply the stacking map $\sigma_2 : (I_2 \circ f_2)(G') \to Y_3^{(u_3(G'))} = Y_3^{(8)}$ as defined above, thereby yielding the map $f_3 = \sigma_2 \circ I_2 \circ f_2 : G' \to Y_3^{(8)}$. Note that σ_2 stacks the 12 many 2-sections of Figure 3 onto S_2^1, with the result shown in Figure 5. The left subfigure shows the stacking of the first 7 sections, while the right subfigure all 12 sections.

Again for each stack address $(x, y) \in S_2^1 \cong \langle Y_2 \rangle$ we indicate by label j, given to various cross sections of this stack, which image sets $\sigma_2(S_2^j)$, $1 \leq j \leq 12$, contribute points to this stack. For example one can check in the right subfigure that for any stack address $(x, 2) \in S_2^1$, $1 \leq x \leq 4$, (so $(x, 2)$ lies in column 2 of S_2^1) the image sets $\sigma_2(S_2^j)$ contributing to $Stack_3((x, 2), 12)$ satisfy $j = 1, 3, 4, 6, 7, 9, 11$ in order of increasing stack height. To see why, refer to Figure 3. There you see that points in column 2 of S_2^3 and S_2^4 "fall through" blank column 2 of S_2^2, points in column 2 of S_2^6 and S_2^7 "fall through" blank columns 2 in S_2^5 and S_2^2, points in column 2 of S_2^9 "fall through" blank columns 2 in S_2^8, S_2^5, and S_2^2, and so on. The resulting stack height is $|Stack_3((x, 2), 12)| = 7$. By contrast, for any stack address $(x, 5) \in S_2^1$, $1 \leq x \leq 4$, the sections contributing to such a stack are $j = 2, 3, 4, 5, 7, 9, 10, 12$ in order of increasing stack height, and the resulting stack height is $|Stack_3((x, 5), 12)| = 8$. In fact the maximum stack height over all 32 stacks addressed by points of S_2^1 is 8.

A more detailed look at the members of various stacks addressed by points of S_2^1 is given in Figure 6. Here we identify the points $x \in G'$ such that $(\sigma_2 \circ I_2 \circ f_2)(x)$ belongs to stack addresses in columns 4 and 5 of S_2^1. For example, consider the members of $Stack_3((3, 4), 7)$ (bolded in the figure), consisting by definition of those images $(\sigma_2 \circ I_2 \circ f_2)(x)$ satisfying $(\sigma_2 \circ I_2 \circ f_2)(x)_{1 \to 2} = (3, 4) \in S_2^1$, and $(I_2 \circ f_2)(x) \in S_2^j$ for some $1 \leq j \leq 7$. Here we must have $(I_2 \circ f_2)(x) = (3, d)$, where $d \equiv 4 (\text{mod } 8)$ by the definition of σ_2 and $8(j - 1) + 1 \leq d \leq 8j$

since $(I_2 \circ f_2)(x) \in S_2^j$. We find such an x when $j = 1$, with $(I_2 \circ f_2)(x) = (3, 4) \in S_2^1$, and Figure 3 shows that $x = (2, 4, 1, 1)$. Here $x_{1\to 3} = (2, 4, 1)$ since x is the 4'th point on chain 2 of D_2^1, while $x_4 = 1$ since $x \in D_3^1$. There is no such x when $j = 2$ since column 4 of S_2^2 is blank. We find such an x when $j = 3$, thus satisfying $(I_2 \circ f_2)(x) = (3, 20) \in S_2^3$, and the figure shows that $x = (2, 4, 3, 1)$ since x is point (2,4) in D_2^3. The remaining two such points x are found similarly; $x = (2, 3, 1, 2) \in D_2^5$ with $(I_2 \circ f_2)(x) = (3, 36) \in S_2^5$, and $x = (2, 4, 2, 2) \in D_2^6$ with $(I_2 \circ f_2)(x) = (3, 44) \in S_2^6$. For the last x note that the first point of D_2^6 in the same chain as x is $(2, 1, 2, 2)$ and $(I_2 \circ f_2)((2, 1, 2, 2)) \in S_2^5$.

To construct $f_4 = \sigma_3 \circ I_3 \circ f_3$, we begin with the parameters needed to define I_3. Observe that $P_3 = 3$ and $e_3 = 7$, so $e_3 - e_2 = 2$ and $S_3^j = P(4) \times P(8) \times P(4)$ for each $1 \le j \le 3$. So I_3 has the form $I_3 : Y_3^{(u_3(G'))} = Y_3^{(8)} \to S_3^{(P_3)} = S_3^{(3)}$. Since $|D_3^j| = 84$, we get by (1) the sequence $s_3(1) = 1$, $s_3(2) = 1$, and $s_3(3) = 2$. A balanced distribution of blank 2-levels (among the sections S_3^j) satisfying (1) together with (5) - (7) is given by the 3×4, $(0, 1)$ matrix $F(3)$ given in Table 1c. We then apply the map $I_3 : Y_3^{(8)} \to S_3^{(3)}$, which distributes the eight 2-levels in $Y_3^{(8)}$ among the twelve 2-levels of $S_3^{(3)}$, leaving four of the latter twelve levels blank.

The result is represented in the left side of Figure 7 as the set of sections S_3^j, $1 \le j \le 3$, each having four 2-levels, with blank 2-levels shaded. For example, S_3^1 has 2-level 1 as blank (as specified by row 1 of $F(3)$), S_3^2 has 2-level 3 as blank (as specified in row 2), and S_3^3 has 2-levels 2 and 4 as blank (as specified in row 3).

Now we apply the stacking map $\sigma_3 : S_3^{(3)} \cap (I_3 \circ f_3)(G') \to Y_4^{(u_4(G'))} = Y_4^{(2)}$ (whose image recall we abbreviate as $\sigma_3(S_3^{(3)})$), which stacks each of the sets $S_3^j \cap (I_3 \circ f_3)(G')$, $1 \le j \le 3$, in succession onto $S_3^1 \cong \langle Y_3 \rangle$ as defined previously. One can check that the maximum stack height $|Stack_4(z, 3)|$ over all 128 stack addresses $z \in \langle Y_3 \rangle$ is 2, again either by checking that each column of $F(3)$ has at most (in fact exactly) 2 zeros, or by seeing in Figure 7 that each of the 128 stack addresses in $\langle Y_3 \rangle$ receives at most 2 points under the map σ_3. In the remainder of Figure 7 we illustrate the stacking map σ_3 in stages. First S_3^2 is stacked on top of $S_3^1 \cong \langle Y_3 \rangle$, with the result that 64 of the stack addresses z (those lying in levels 2 and 4 of $\langle Y_3 \rangle$) so far have 4-dimensional stack height 2 (i.e. $|Stack_4(z, 2)| = 2$ for these z), while 64 of the stack addresses z (those lying in levels 1 and 3 of $\langle Y_3 \rangle$) so far have 4-dimensional stack height 1 (i.e. $|Stack_4(z, 2)| = 1$ for these z). Next, S_3^3 is stacked on top of the previous stacking, so that now every stack address has 4-dimensional stack height 2 (i.e. $|Stack_4(z, 3)| = 2$ for all $z \in \langle Y_3 \rangle$).

Consider the final result $\sigma_3(S_3^{(3)})$, yielding the map f_4, illustrated in the upper right of Figure 7. There, focus on the stack addresses $(x, y, z) \in \langle Y_3 \rangle$ with $z = 2$, $1 \le x \le 4$, $1 \le y \le 8$, these lying in the second 2-level of $\langle Y_3 \rangle$ (since $z = 2$). The sets $\sigma_3(S_3^j)$, $1 \le j \le 3$, contributing points to any such stack $Stack_4((x, y, 2), 3)$ satisfy $j = 1$ or 2, where in Figure 7 the contribution of $\sigma_3(S_3^1)$ is represented by $1b$, and of $\sigma_3(S_3^2)$ by $2b$.

In Figure 8 we look in detail at the individual stacks $Stack_4((3, 1, 2), 3)$, $Stack_4((3, 4, 2), 3)$, and $Stack_4((1, 4, 2), 3)$. For example, the members of $Stack_4((3, 1, 2), 3)$ are the two images $(\sigma_3 \circ I_3 \circ f_3)(\alpha)$, indicated by their preimages $\alpha \in G'$ which are $\alpha = (3, 2, 2, 1)$ and $\alpha = (2, 1, 4, 2)$ in increasing order of stack height.

4 Tools for the general construction

In this section we develop two tools used in our general construction:
(1) the designation of blank $(i-1)$-levels in i-sections S_i^j, $1 \le j \le P_i$, and
(2) the construction of an ordering (by consecutive integer labels) of the vertices of any hypercube such that for any reasonably long interval of successive label values, any two vertices whose labels lie in that interval are at fairly small hypercube distance.

4.1 The Construction of Blank Levels

In this subsection we describe the sequence $\{s_i(j)\}$, $1 \le i \le k$, $1 \le j \le P_i$, where $s_i(j)$ is the number of $(i-1)$-levels of S_i^j that are designated blank under the map $I_i \circ f_i$. We show that this sequence satisfies equation (1). That is, for each $1 \le r \le P_i$ we show that $S_i^{(r)}$ has just enough nonblank $(i-1)$-levels to contain $(I_i \circ f_i)(D_i^{(r)})$. Finally, we construct for each j the actual set of $s_i(j)$ many $(i-1)$-levels in S_i^j that are designated blank, and show that the required properties (5)-(7) are satisfied. This construction is based on a theorem of Knuth on simultaneous roundings of sequences.

Given $G = [a_1 \times a_2 \times \cdots \times a_k]$ and $1 \le i \le k-1$, define the sequence $\{s_i(j)\}$, with $1 \le j \le P_i$, by

$$s_i(j) = 2^{e_i - e_{i-1}} - \lceil \frac{a_1 a_2 \cdots a_i}{2^{e_{i-1}}} \rceil + \lfloor j\phi_i \rfloor - \lfloor (j-1)\phi_i \rfloor \tag{8}$$

where $\phi_i = \lceil \frac{a_1 a_2 \cdots a_i}{2^{e_i - 1}} \rceil - \frac{a_1 a_2 \cdots a_i}{2^{e_i - 1}}$.

Lemma 4.1 *Let $P_i = a_{i+1} a_{i+2} \cdots a_k$, $1 \le r \le P_i$, and $1 \le i \le k$.*
(a) The sequence $\{s_i(j)\}$, $1 \le j \le P_i$, defined above satisfies $\lceil \frac{r a_1 a_2 \ldots a_i}{2^{e_i - 1}} \rceil + \sum_{j=1}^{r} s_i(j) = r 2^{e_i - e_{i-1}}$. In particular, taking $s_i(j)$ to be the number of blank $(i-1)$-levels of S_i^j, then the number of nonblank $(i-1)$-levels in $S_i^{(r)}$ is $\lceil \frac{r a_1 a_2 \ldots a_i}{2^{e_i - 1}} \rceil = \lceil \frac{|D_i^{(r)}|}{2^{e_i - 1}} \rceil$.
(b) $s_i(j) = p$ or $p+1$, where $p = 2^{e_i - e_{i-1}} - \lceil \frac{a_1 a_2 \cdots a_i}{2^{e_i - 1}} \rceil$. Also $\frac{s_i(j)}{2^{e_i - e_{i-1}}} \le \frac{1}{2}$.

Proof. For (a), observe that the sum $\sum_{j=1}^{r} s_i(j)$ telescopes. Note also that $r \lceil \frac{a_1 a_2 \ldots a_i}{2^{e_i - 1}} \rceil$ is an integer. Thus $\sum_{j=1}^{r} s_i(j) = r 2^{e_i - e_{i-1}} - r \lceil \frac{a_1 a_2 \ldots a_i}{2^{e_i - 1}} \rceil + \lfloor r \lceil \frac{a_1 a_2 \ldots a_i}{2^{e_i - 1}} \rceil - r \frac{a_1 a_2 \ldots a_i}{2^{e_i - 1}} \rfloor = r 2^{e_i - e_{i-1}} + \lfloor -r \frac{a_1 a_2 \ldots a_i}{2^{e_i - 1}} \rfloor = r 2^{e_i - e_{i-1}} - \lceil r \frac{a_1 a_2 \ldots a_i}{2^{e_i - 1}} \rceil$. Part (a) follows.

Consider (b). Since $\phi_i < 1$ we have $0 \le \lfloor j\phi_i \rfloor - \lfloor (j-1)\phi_i \rfloor \le 1$, giving the first statement. Since $a_1 a_2 \cdots a_i \ge 2^{e_i - 1} + 1$, we get $s_i(j) \le 2^{e_i - e_{i-1}} - \lceil \frac{2^{e_i - 1} + 1}{2^{e_i - 1}} \rceil + 1 \le 2^{e_i - e_{i-1} - 1}$. The second statement follows. ∎

We now specify, for each $2 \le i \le k-1$ and $1 \le j \le P_i$, which $s_i(j)$ out of the $2^{e_i - e_{i-1}}$ many $(i-1)$-levels in S_i^j will be designated blank. Keep in mind that this designation must satisfy the balance properties (5)-(7) we required in the overview. For this, we need the following theorem of Knuth [19].

Let $x_1, \ldots, x_n$ be a sequence of reals, and γ a permutation of $\{1, 2, \ldots, n\}$. Let $S_k = x_1 + \cdots + x_k$ and $\Sigma_k = x_{\gamma(1)} + \cdots + x_{\gamma(k)}$ be the partial sums for these two independent orderings of the x_i's. Consider a *rounding* of the x_i's; that is, a designation of integers $\bar{x}_i$ satisfying $\lfloor x_i \rfloor \le \bar{x}_i \le \lceil x_i \rceil$ for $1 \le i \le n$. Now let the corresponding partial sums be

$\bar{S}_k = \bar{x}_1 + \cdots + \bar{x}_k$ and $\overline{\Sigma}_k = \bar{x}_{\gamma(1)} + \cdots + \bar{x}_{\gamma(k)}$. We say that a rounding of the x_i's is *consistent* with the original sequence $\{x_i\}$ (resp. with the permuted sequence under γ) if $\lfloor S_k \rfloor \leq \bar{S}_k \leq \lceil S_k \rceil$ (resp. $\lfloor \Sigma_k \rfloor \leq \overline{\Sigma}_k \leq \lceil \Sigma_k \rceil$) for each $1 \leq k \leq n$. We also say that the rounding is a *two-way* rounding if it is simultaneously consistent with both the original sequence and the permuted sequence under γ.

Theorem 4.2 *[19] For any finite sequence $x_1, \ldots, x_n$ of reals and any permutation γ of $\{1, 2, \ldots, n\}$, there is a two-way rounding of the x_i.*

The existence of two-way roundings was shown earlier by Spencer [27] by probabilistic methods, as a corollary to more general results on the discrepancy of set systems [23]. Knuth's network flow based proof of Theorem 4.2, omitted here, is constructive and yields improved error bounds. The two-way rounding produced is not necessarily unique.

As a consequence of Knuth's theorem, we obtain the following theorem (from [11] and [12]) on roundings of matrices which are consistent with respect to all initial row and column sums. We give its short proof for completeness. This extends the rounding lemma of Baranyai ([4]) giving such consistency with respect to all row sums, to all column sums, and to the sum of all matrix entries. Additional results on roundings of matrices, including extensions of previous work, applications to digital halftoning, and improved running time and error bounds in implementation can be found in the work of Doerr ([10], [11], [12] and others), Asano ([1] and [2]), and Wright [29] among others.

Theorem 4.3 *Let $T = (t_{ij})$ be an $m \times n$ matrix with $0 \leq t_{ij} \leq 1$ for all i and j. Then there exists an $m \times n$, $(0, 1)$ "rounding" matrix $F = (f_{ij})$ of T; that is, where $f_{ij} = \lfloor t_{ij} \rfloor$ or $\lceil t_{ij} \rceil$, satisfying the following properties.*

(a) *For each b and i, $1 \leq b \leq n$, $1 \leq i \leq m$, we have $|\sum_{j=1}^{b}(t_{ij} - f_{ij})| < 1$.*

(b) *For each b and j, $1 \leq b \leq m$, $1 \leq j \leq n$, we have $|\sum_{i=1}^{b}(t_{ij} - f_{ij})| < 1$.*

(c) $|\sum_{i=1}^{m}\sum_{j=1}^{n}(t_{ij} - f_{ij})| < 1$.

Proof. For the most part we paraphrase proofs in [11] and [12]. First consider parts (a) and (b). We construct an $(m+1) \times (n+1)$ matrix Y from T which has integral row and column sums by appending to each row and each column a last entry in $[0, 1)$ just large enough to make that row or column have an integer sum. Specifically, let $y_{ij} = t_{ij}$ for $1 \leq i \leq m$ and $1 \leq j \leq n$, and then $y_{m+1,j} = \lceil \sum_{i=1}^{m} t_{ij} \rceil - \sum_{i=1}^{m} t_{ij}$ for $1 \leq j \leq n$, and $y_{i,n+1} = \lceil \sum_{j=1}^{n} t_{ij} \rceil - \sum_{j=1}^{n} t_{ij}$ for $1 \leq i \leq m$, and $y_{m+1,n+1} = \sum_{i=1}^{m}\sum_{j=1}^{n} t_{ij}$. Then Y has integral row and column sums.

Consider now the following two orderings of the entries of Y. First we order these entries by "row-major" order; that is, first the entries of row 1, then those of row 2, etc., until row $m+1$, and within any row i place y_{ij} ahead of y_{ir} iff $j < r$. Similarly consider the "column-major" order where we place y_{ij} before y_{rs} iff either $j < s$, or $j = s$ and $i < r$.

Applying Knuth's theorem, there is a (not necessarily unique) two-way rounding matrix $\overline{Y} = \bar{y}_{ij}$ relative to these two orders. Since every row and column sum of Y is already an integer, we get $\sum_{i=1}^{p}\sum_{j=1}^{n+1}(y_{ij} - \bar{y}_{ij}) = 0$ for any $1 \leq p \leq m+1$, and $\sum_{j=1}^{q}\sum_{i=1}^{m+1}(y_{ij} - \bar{y}_{ij}) = 0$ for any $1 \leq q \leq n+1$. Thus taking suitable partial sums in the row major order we get initial row sum estimates $|\sum_{j=1}^{b}(y_{ij} - \bar{y}_{ij})| < 1$ for any fixed i and $1 \leq b \leq n+1$. Similarly, suitable partial

sums in column major order yield $|\sum_{i=1}^{b}(y_{ij}-\bar{y}_{ij})|<1$ for any fixed j and $1\le b\le m+1$. Finally, let F be the upper left $m\times n$ submatrix of $\overline{Y}$; that is, $F=(f_{ij})$ where $f_{ij}=\bar{y}_{ij}$ for $1\le i\le m$ and $1\le j\le n$. Then recalling that $t_{ij}=y_{ij}$ for such i and j, we get parts (a) and (b) of the theorem.

For (c), let $R=\sum_{i=1}^{m+1}\sum_{j=1}^{n+1}(y_{ij}-\bar{y}_{ij})$, $S=\sum_{i=1}^{m+1}(y_{i,n+1}-\bar{y}_{i,n+1})$, and $S'=\sum_{j=1}^{n+1}(y_{m+1,j}-\bar{y}_{m+1,j})$, noting that these three quantities are all 0 since each row or column sum in Y is integral. Then $|\sum_{i=1}^{m}\sum_{j=1}^{n}(t_{ij}-f_{ij})|=|R-S-S'+(y_{m+1,n+1}-\bar{y}_{m+1,n+1})|\le|y_{m+1,n+1}-\bar{y}_{m+1,n+1}|<1$. ∎

The setting in which we apply Theorem 4.3 is as follows. Let $X=\{s_1,s_2,\ldots,s_m\}$ be an integer sequence such that for all $1\le i\le m$ we have $s_i=k$ or $k+1$ for some fixed integer k independent of i. Also let n be a positive integer satisfying $k+1\le n$. Define the $m\times n$ matrix $T^X=(t_{ij})$ as follows. For each fixed row index i, $1\le i\le m$, let $t_{ij}=\frac{s_i}{n}$ for all j, $1\le j\le n$. Thus all entries in a given row of T^X have the same constant value. For some rows this constant is $\frac{k}{n}$ while for the rest it is $\frac{k+1}{n}$, and row i of T^X has row sum s_i. Now let $F^X=(f_{ij})$ be an $m\times n$, $(0,1)$ rounding matrix of T^X as guaranteed to exist by Theorem 4.3; that is, with T^X and F^X playing the roles of T and F respectively in that theorem. For suitably chosen integers m,n and integer sequence X determined by the construction in the next section, this F^X will be, for fixed i, the matrix $F(i)$ introduced in the previous section that encodes which $(i-1)$-levels of Y_i will be designated blank. This F^X has certain balance properties described in the following theorem.

Theorem 4.4 *Let k and n be positive integers with $1\le k+1\le n$, and $X=(s_1,s_2,\cdots,s_m)$ a sequence with $s_i=k$ or $k+1$ for all i. Let $T^X=(t_{ij})$ and $F^X=(f_{ij})$, a rounding of T^X, be the $m\times n$ matrices as defined above. Then F^X has the following properties.*

(a) $\sum_{j=1}^{n}f_{ij}=s_i$ *for* $1\le i\le m$.

(b) $|\sum_{j=1}^{b}f_{jr}-\sum_{j=1}^{b}f_{js}|\le 1$ *for* $1\le r,s\le n$, $1\le b\le m$.

(c) $|\sum_{j=1}^{b}f_{rj}-\sum_{j=1}^{b}f_{sj}|\le 2$ *for* $1\le r,s\le m$, $1\le b\le n$.

Proof. For (a), note that row i of T^X has the integer s_i as its sum. Then (a) follows from Theorem 4.3a on taking $b=n$.

For (b), observe that for any $b, 1\le b\le m$, the sequences formed by the first b entries in any two columns r and s of T^X are identical. So $\sum_{j=1}^{b}t_{jr}=\sum_{j=1}^{b}t_{js}$ for any r and s. Thus by Theorem 4.3b, $\sum_{j=1}^{b}f_{jr}$ and $\sum_{j=1}^{b}f_{js}$ are roundings of the same quantity, and hence can differ by at most 1, proving (b).

For (c), recall that any two rows r and s of T^X are either identical, or one has constant entries $\frac{k}{n}$ and the other constant entries $\frac{k+1}{n}$. So the maximum difference between any two initial row sums of T^X occurs in the second possibility, and in that case the difference we get is $|\sum_{j=1}^{b}t_{rj}-\sum_{j=1}^{b}t_{sj}|\le\frac{b}{n}\le 1$. By Theorem 4.3c the initial row sums $\sum_{j=1}^{b}f_{rj}$ and $\sum_{j=1}^{b}f_{sj}$ of F^X are roundings of the corresponding initial row sums in T^X. Since the latter sums differ by at most 1, part (c) follows, where the upper bound of 2 can be achieved only if there is an integer strictly between the corresponding initial row sums in T^X. ∎

For any two integers r and s, $1\le r,s\le m$, we will need to bound the column difference between the p'th zero from the left in row r of F^X and the q'th zero from the left in row s of

F^X as a function of $|p-q|$, independent of r and s. For this purpose, let $N_i(d)$ be the column index of the d'th zero from the left in row i of F^X; that is, $N_i(d) = min\{b : b = d + \sum_{j=1}^{b} f_{ij}\}$. We omit mention of X in the notation $N_i(d)$, since X will be clear by context. We will call the entry f_{ih} *forward* if $\sum_{j=1}^{h} f_{ij} = \lceil \sum_{j=1}^{h} t_{ij} \rceil$. Otherwise we call f_{ih} *backward*, so in that case $\sum_{j=1}^{h} f_{ij} = \lceil \sum_{j=1}^{h} t_{ij} \rceil - 1$. Recall that every entry of F^X is either forward or backward by Theorem 4.3a.

It will be convenient to interpret the entries f_{ij} of the $m \times n$ matrix F^X and the function $N_i(*)$ using "wraparound". For example we let $f_{i,n+3} = f_{i+1,3}$. Similarly if row i of F^X has q many 0's then we let $N_i(q+3) = N_{i+1}(3)$, while if row $i+1$ has q' many 0's then let $N_i(q+q'+3) = N_{i+2}(3)$, and so on. Also for $t \leq 0$, let $N_{i+1}(t) = N_i(q+t)$.

Corollary 4.5 *Let $X = \{s_1, s_2, \ldots, s_m\}$, $T^X = (t_{ij})$, $F^X = (f_{uv})$, and k be as in the preceding theorem, and assume now that $k+1 \leq \frac{n}{2}$. Let $c = \frac{k+1}{n}$. Then the $m \times n$, $(0,1)$-matrix F^X has the following properties.*

(a) *If f_{ih} is forward, then $\sum_{j=h+1}^{h+2e} f_{ij} \leq e$. In particular, if $f_{i,N_i(d)}$ is forward, then $N_i(d+e) - N_i(d) \leq 2e$.*

(b) *If f_{ih} is backward, then $\sum_{j=h+1}^{h+2e} f_{ij} \leq e+1$. In particular, if $f_{i,N_i(d)}$ is backward, then $N_i(d+e) - N_i(d) \leq 2e+2$.*

(c) *For integers r, s, d, e with $1 \leq r, s \leq m$, we have $N_r(d+e) - N_s(d) \leq 2e+4$.*

(d) *Let $2 \leq i \leq k-1$. Set $m = P_i = a_{i+1}a_{i+2}\cdots a_k$, $n = 2^{e_i - e_{i-1}}$, $k = min\{s_i(j) : 1 \leq j \leq P_i\}$, $s_j = s_i(j)$, $1 \leq j \leq m = P_i$, and $f_{uv} = f_{uv}(i)$. With these settings, F^X satisfies (5)-(7) (which include the conditions (2)-(4) as the special case $i = 2$).*

Proof. For part (a), begin by observing that for any positive integer p we have $\sum_{j=N_i(d)+1}^{N_i(d)+p} t_{ij} \leq pc \leq \frac{p}{2}$. Now take $p = 2e$ to be an even integer. Then since f_{ih} is forward, the number of 1's among the entries f_{ij}, $h+1 \leq j \leq h+2e$, is at most e, so $\sum_{j=h+1}^{h+2e} f_{ij} \leq e$ as claimed. For the second statement, since the number of these entries is $2e$ and $\sum_{j=h+1}^{h+2e} f_{ij} \leq e$, it follows that the number of 0's among these must entries is at least e, and hence $N_i(d+e) - N_i(d) \leq 2e$.

For part (b), we proceed as in part (a), taking $p = 2e$ to be an even integer. The difference is that since $f_{i,N_i(d)}$ is backward, the number of 1's among the entries f_{ij}, $h+1 \leq j \leq h+2e$ can be at most $1+e$ (since that number now includes the 1 corresponding to $\lceil \sum_{j=1}^{N_i(d)} t_{ij} \rceil$). It follows that $\sum_{j=h+1}^{h+2e} f_{ij} \leq e+1$. For the second statement, since $\sum_{j=h+1}^{h+2e} f_{ij} \leq e+1$ it follows that the number of 0's among these entries is at least $e-1$. Now replacing $2e$ by $2e+2$ in the above reasoning, we find that the number of 0's among the $2e+2$ entries f_{ij}, $h+1 \leq j \leq h+2e+2$, is at least e. It follows that $N_i(d+e) - N_i(d) \leq 2e+2$.

Consider now part (c). By Theorem 4.4c we have $N_r(d-2) \leq N_s(d)$; that is, there are at least $d-2$ many 0's among the entries f_{rj}, $1 \leq j \leq N_s(d)$. Suppose first that $N_r(d-1) > N_s(d)$, so that there are exactly $d-2$ many 0's among the entries f_{rj}, $1 \leq j \leq N_s(d)$. It follows that $\sum_{j=1}^{N_s(d)} f_{rj} - \sum_{j=1}^{N_s(d)} f_{sj} = 2$. Thus $f_{r,N_s(d)}$ is forward. So by part (a) we have $\sum_{N_s(d)+1}^{N_s(d)+2e+4} f_{rj} \leq e+2$. So there are at least $e+2$ many 0's among the entries f_{rj}, $N_s(d)+1 \leq j \leq N_s(d)+2e+4$. Combining these 0's with the $d-2$ many 0's among the entries f_{rj}, $1 \leq j \leq N_s(d)$, obtain $N_r(d+e) \leq N_s(d) + 2e + 4$, as required. Now assume $N_r(d-1) \leq N_s(d)$. Then we have

$N_r(d+e) - N_s(d) \leq N_r(d+e) - N_r(d-1) \leq 2(e+1) + 2 = 2e+4$, where we have used the more generous bound from part (b) in bounding $N_r(d+e) - N_r(d-1)$. This completes (c).

Finally consider part (d). By Lemma 4.1b we have $s_i(j) = k$ or $k+1$ for each term $s_i(j)$, $1 \leq j \leq P_i$, of our sequence X. Hence we may apply Theorem 4.4a,b,c with the given settings for m, n, X, and for the entries f_{uv} of F^X, to obtain (5), (6), and (7) respectively. ■

We now apply Theorems 4.3 and 4.4 to construct the matrices $F(i)$ of the previous section. Recall that $F(i) = (f_{cd}(i))$ is the $P_i \times 2^{e_i - e_{i-1}}$, (0,1)-matrix such that for $1 \leq d \leq 2^{e_i - e_{i-1}}$ and $1 \leq c \leq P_i$, the $(i-1)$-level $Y_i^{(c-1)2^{e_i - e_{i-1}} + d}$ of i-section S_i^c (i.e. the d'th $(i-1)$-level of S_i^c) is blank precisely when $f_{cd}(i) = 1$.

Construction of the matrix $F(i)$, $1 \leq i \leq k-1$

1. Given G and fixed i, compute the sequence $X = \{s_i(j)\}, 1 \leq j \leq P_i$, using formula (8).

2. Form the $P_i \times 2^{e_i - e_{i-1}}$ matrix T^X and construct rounding F^X of T^X as in Theorem 4.4; that is, letting $m = P_i$, $n = 2^{e_i - e_{i-1}}$, and $s_j = s_i(j)$ in the definition of T^X (preceding Theorem 4.4). We note that the hypotheses of Theorem 4.4 hold with this X by Lemma 4.1b.

3. Let $F(i) = F^X$.

To illustrate, recall $G'' = [3 \times 7 \times 4]$ from the previous section. Toward constructing $F(2)$, apply formula (8) to obtain the sequence $X = \{s_2(1) = 2, s_2(2) = 3, s_2(3) = 3, s_2(4) = 3\}$. So the $P_2 \times 2^{e_2 - e_1} = 4 \times 8$ matrix T^X has constant row entries $\frac{s_2(r)}{8} = \frac{1}{4}$ or constant row entries $\frac{s_2(r)}{8} = \frac{3}{8}$ in any given row r, $1 \leq r \leq 4$, as shown at left in Table 1a. The rounding of T^X by Knuth's network flow method or Doerr's approach (Theorems 4.2 and 4.3) yields, as one possibility, the $(0,1)$ matrix $F(2) := F^X$ at right in Table 1a. Next recall $G' = [3 \times 7 \times 4 \times 3]$. To construct $F(2)$, apply formula (8) to obtain the sequence $X = \{s_2(j)\}$, $1 \leq j \leq 12$, given by $X = (2, 3, 3, 3, 2, 3, 3, 3, 2, 3, 3, 3)$. So the $P_2 \times 2^{e_2 - e_1} = 12 \times 8$ matrix T^X has constant row entries $\frac{1}{4}$ or $\frac{3}{8}$ in any given row, and is shown at left in Table 1b. Again applying Theorems 4.2 and 4.3 gives a possible rounding of T^X given by $F(2) = F^X$ at right in Table 1b. Toward constructing $F(3)$, still for G', we apply formula (8) to obtain the sequence $X = \{s_3(1) = 1, s_3(2) = 1, s_3(3) = 2\}$. The $P_3 \times 2^{e_3 - e_2} = 3 \times 4$ matrix T^X therefore has constant row entries $\frac{1}{4}$ or $\frac{1}{2}$, as shown at left in Table 1c, while a possible rounding $F(3) = F^X$ by these theorems is shown at right.

4.2 An Integer Labeling of Hypercubes

In this subsection we construct an ordering L_t of the vertices of Q_t such that for any interval of at most $O(log(t))$ consecutive points under this ordering, any two points x, y in that interval satisfy $dist_{Q_t}(x, y) \leq 3$. This labeling is based on the existence of certain spanning subgraphs of hypercubes of suitable dimension, as follows. Define a *cyclic caterpillar* as a connected graph H such that removal of all leaves of H results in a cycle graph C_e for some $e \geq 3$. A cyclic caterpillar is *r-regular* if each vertex of its cycle subgraph C_e has exactly r neighboring leaves not on the cycle. Denote such an r-regular cyclic caterpillar by $Cat(e, r)$.

We are interested in finding spanning subgraphs $Cat(e, r)$ of hypercubes. Clearly if $Cat(e, r)$ spans Q_n, then $e = \frac{2^n}{r+1}$, so $r+1$ must be a power of 2. Papers on this subject include [7],

[13], and [17], while [3] examines the general question of finding short dominating cycles or paths in the hypercube. We use following result.

Theorem 4.6 *[Corollary 5.8 in [7]] There exists a spanning cyclic regular caterpillar* $Cat(e, 2r+1)$ *of* Q_n *provided that* $r+1 = 2^i$ *and* $n = 2^{i+1} + 2i$ *for some integer* $i \geq 0$.

From this we easily obtain the following.

Corollary 4.7 *Suppose* $r+1 = 2^i$ *and* $t \geq 2^{i+1} + 2i$ *for an integer* $i \geq 0$. *Then* Q_t *contains a spanning cyclic* $(2r+1)$*-regular caterpillar* $Cat(e, 2r+1)$ *for suitable* e.

Proof. Recall that Q_{t+1} is the cartesian product $Q_{t+1} = Q_t \times K_2$. Hence if Q_t contains a spanning subgraph $Cat(e, 2r+1)$, then Q_{t+1} contains a spanning subgraph $Cat(2e, 2r+1)$. The corollary follows by induction on t. ■

We can now construct our desired labeling of hypercubes of sufficiently large dimension.

Corollary 4.8 (a) *Let* r, i, t *be positive integers satisfying* $r+1 = 2^i$ *and* $t \geq 2^{i+1} + 2i$. *Then there exists a one to one integer labeling* $L_t : V(Q_t) \to \{1, 2, \ldots, 2^t\}$ *such that for any* $x, y \in V(Q_t)$ *we have* $|L_t(x) - L_t(y)| \leq 2r+3 \Rightarrow dist_{Q_t}(x, y) \leq 3$, *where the indicated difference is taken modulo* 2^t.
(b) *Let* $t \geq 22$. *Then there exists a one to one integer labeling* $L_t : V(Q_t) \to \{1, 2, \ldots, 2^t\}$ *such that for any* $x, y \in V(Q_t)$ *we have* $|L_t(x) - L_t(y)| \leq 17 \Rightarrow dist_{Q_t}(x, y) \leq 3$, *where the indicated difference is taken modulo* 2^t.

Proof. For (a), consider the spanning subgraph $Cat(e, 2r+1)$ of Q_t, $t \geq 2^{i+1} + 2i$, from Corollary 4.7. For $1 \leq i \leq e$, let x_i be the vertices of the cycle C_e in $Cat(e, 2r+1)$ indexed consecutively around this cycle. Also let $x_{i,1}, x_{i,2}, \ldots, x_{i,2r+1}$ be the leaf neighbors of x_i. Now define L_t by letting $L_t(x_i) = (2r+2)i$ for $1 \leq i \leq e$, and $L_t(x_{i,j}) = (2r+2)(i-1) + j$ for fixed i and $1 \leq j \leq 2r+1$. For the claim, the critical case to check is when x (or y) $= x_{i,2r+1}$ for any $1 \leq i \leq e$. The $2r+3$ points which follow this x in this ordering are (in order) x_i, $\{x_{i+1,j} : 1 \leq j \leq 2r+1\}$, and x_{i+1}, all at distance 3 from x. Further $2r+3$ is best possible since the $(2r+4)$'th point following x is $x_{i+2,1}$, and $dist_{Q_t}(x, x_{i+2,1}) = 4$.
For (b), simply apply part (a) with $r = 7$, yielding $i = 3$, and $t \geq 22$. ■

We will apply the labeling L_t when $t = e_i - e_{i-1}$, for each $1 \leq i \leq k$. So to apply Corollary 4.8b we need the condition $t = e_i - e_{i-1} \geq 22$. We therefore assume $a_i \geq 2^{22}$, $1 \leq i \leq k$, from now on, which ensures that this condition holds.

5 The general construction

In this section we inductively construct a series of embeddings $f_i : G \to [\langle Y_{i-1} \rangle \times P(u_i)] = Y_i^{(u_i)} \subset Y_i$, $3 \leq i \leq k$, where $u_i = \lceil \frac{|G|}{|\langle Y_{i-1} \rangle|} \rceil = \lceil \frac{|G|}{2^{e_{i-1}}} \rceil$. At the end we relabel the points of $\langle Y_{k-1} \rangle \times P(u_k)$ with hypercube addresses coming from $Opt(G)$, using the inverse of the labeling of Corollary 4.8. The composition of this relabeling with the map f_k yields the final embedding $H^k : G \to Opt(G)$. We follow the general plan outlined in section 3.

Recall $S_i(G)$, the set of points of $S_i^{(P_i)}$ lying in nonblank columns of $S_i^{(P_i)}$. We let $S_i^r(G) = S_i(G) \cap S_i^r$, the set of points lying in nonblank columns of S_i^r, and $S_i^{(r)}(G) = \bigcup_{j=1}^r S_i^j(G)$. Recall we stipulated that $(I_i \circ f_i)(G) \subseteq S_i(G)$. For $r \leq P_i$ we let $S_i^r(G)' = S_i^r(G) \cap (I_i \circ f_i)(G)$ and $S_i^{(r)}(G)' = S_i^{(r)}(G) \cap (I_i \circ f_i)(G)$. Thus $S_i^{(P_i)}(G)'$ is the domain of the stacking map σ_i, and $S_i^r(G)'$ (resp. $S_i^{(r)}(G)'$) is the subset of that domain lying in section S_i^r (resp. $S_i^{(r)}$). We will see later that $S_i^{(P_i-1)}(G) = S_i^{(P_i-1)}(G)'$.

5.1 The three dimensional embedding

In this subsection we construct the map $f_3 : G \to [\langle Y_2 \rangle \times P(u_3)] \subset Y_3$ for a 3-dimensional grid $G = [a_1 \times a_2 \times a_3]$. Since $f_3(G)$ is a 3-dimensional grid, we can visualize its construction along lines of Section 3 using Figures 3, 4, and 5, to which we refer the reader toward following the construction which follows. This 3-dimensional case will hopefully help in understanding the generalization to higher dimensions in the next subsection.

Construction of the map $f_3 : [a_1 \times a_2 \times a_3 \times \cdots \times a_k] \to Y_3$

1. Set $G = [a_1 \times a_2 \times a_3 \times \cdots \times a_k]$. Construct the embedding $f_2 : G \to Y_2^{(u_2)}$ of section 2.

2. [**Designation of Blank Columns**]

a) Set $P_2 = \Pi_{i=3}^k a_i$.

b) Construct the matrix $F(2) = (f_{ij}(2))$ by the procedure following Corollary 4.5 for the case $i = 2$. (Comment: The matrix entries $f_{ij}(2)$ now satisfy (2)-(4) by Corollary 4.5d.)

c) Now designate 'blank' columns in the subgraph $S_2^{(P_2)}$ of Y_2 as follows.

Column j of S_2^i is 'blank' if $f_{ij}(2) = 1$, and is 'nonblank' if $f_{ij}(2) = 0$.

(Comment: Recall that column j of S_2^i is column $(i-1)2^{e_2-e_1} + j$ of Y_2.)

d) For $1 \leq r \leq P_2$, let $S_2^r(G)$ (resp. $S_2^{(r)}(G)$) be the set points lying in nonblank columns of S_2^r (resp. $S_2^{(r)}$).

3. [**The Inflation Step**]

a) "Inflate" the image $f_2(G)$ by the map $I_2 : Y_2^{(u_2(G))} \to S_2^{(P_2)}(G)$ given as follows. For any $(a, b) \in Y_2^{(u_2(G))}$, let

$$I_2(a, b) = (a, b'),$$

where $Y_2^{b'}$ is the b'th nonblank column in $S_2^{(P_2)}$ in increasing order of b.

b) Let $S_2^r(G)' = S_2^r(G) \cap (I_2 \circ f_2)(G)$ and $S_2^{(r)}(G)' = S_2^{(r)}(G) \cap (I_2 \circ f_2)(G)$.

4. [**The Stacking Step**]

"Stack" the sets $S_2^r(G)'$, $1 \leq r \leq P_2$, successively "over" $S_2^1 \cong \langle Y_2 \rangle$ by the stacking map $\sigma_2 : S_2^{(P_2)}(G)' \to S_2^1 \times P(u_3) = Y_3^{(u_3)}$ defined as follows. Take $y = (a, b) \in S_2^{(P_2)}(G)'$, say with $y \in S_2^r(G)'$.

a) Let $n_y = |\{z \in S_2^{(r)}(G)' : z_1 = a \text{ and } z_2 \equiv b \pmod{2^{e_2-e_1}}\}|$.

b) Let $\bar{b}$ satisfy $b \equiv \bar{b} \pmod{2^{e_2-e_1}}$, with $1 \leq \bar{b} \leq 2^{e_2-e_1}$.

c) Define $\sigma_2(a, b) = (a, \bar{b}, n_y)$.

(Comment: Take any $(a,b) \in S_2(G)'$. Then $1 \leq \sigma_2(a,b)_2 = \bar{b} \leq 2^{e_2-e_1}$, and $1 \leq \sigma_2(a,b)_1 = a \leq 2^{e_1}$. Thus $\sigma_2(a,b)_{1\to 2} \in S_2^1$ so we can view σ_2 as "stacking" the sets $S_2^r(G)', 1 \leq r \leq P_2$, in succession by increasing r "over" $S_2^1 \cong \langle Y_2 \rangle$. So we have $(\sigma_2 \circ I_2 \circ f_2)(G) \subseteq \langle Y_2 \rangle \times P(u) \subset Y_3$, where u is the maximum of n_y over all y belonging to the last set $S_2^{P_2}(G)'$. We will prove later that $u = u_3(G)$.)

5. [**The Composition Step**]
Finally, define $f_3 : G \to Y_3$ as the composition $f_3 = \sigma_2 \circ I_2 \circ f_2$.

5.2 Embeddings of grids of higher dimension

Again set $G = [a_1 \times \cdots \times a_k]$. In this subsection we inductively construct embeddings $f_i : G \to Y_i^{(u_i)}$ for $3 \leq j \leq k$. Assume then that we have constructed the required maps $f_2, \ldots, f_i$, $3 \leq i < k$, and we construct $f_{i+1} : G \to Y_{i+1}^{(u_{i+1})}$.

Starting from $f_i(G) \subset Y_i^{(u_i)}$, we will use direct analogues I_i and σ_i of the inflation map I_2 and the stacking map σ_2 used in constructing f_3 from f_2. In particular, I_i will inflate $f_i(G)$ by introducing blank $(i-1)$-levels Y_i^j of Y_i using a matrix $F(i) = F^X$ (constructed by the procedure following Corollary 4.5) that encodes which $(i-1)$-levels of Y_i will be introduced as blank. Then a stacking map $\sigma_i : S_i^{(P_i)}(G)' \to S_i^1 \times P(u_{i+1}) = Y_{i+1}^{(u_{i+1})}$ stacks the sets $S_i^r(G)'$, $1 \leq r \leq P_i$, in succession by increasing r over $S_i^1 \cong \langle Y_i \rangle$ to yield the final image $f_{i+1}(G)$. Thus we may write f_{i+1} as the composition $f_{i+1} = \sigma_i \circ I_i \circ f_i$, and inductively $f_{i+1} = \sigma_i \circ I_i \circ \sigma_{i-1} \circ I_{i-1} \circ \cdots \circ \sigma_2 \circ I_2 \circ f_2$.

Construction of the map $f_{i+1} : [a_1 \times a_2 \times a_3 \times \cdots \times a_k] \to Y_{i+1}^{(u_{i+1})}$, $2 \leq i < k$

1. Set $G = [a_1 \times a_2 \times a_3 \times \cdots \times a_k]$. Assume inductively that the map $f_i : [a_1 \times a_2 \times a_3 \times \cdots \times a_k] \to Y_i^{(u_i)}$ has been constructed. We now operate on $f_i(G)$ to obtain our image $f_{i+1}(G) \subset Y_{i+1}^{(u_{i+1})}$.

2. [**Designation of Blank** $(i-1)$**-Levels**]
a) Set $P_i = \Pi_{t=i+1}^k a_t$.
b) Construct the $P_i \times 2^{e_i - e_{i-1}}$, $(0,1)$-matrix $F(i) = (f_{cd}(i))$ by the procedure following Corollary 4.5.
[Comment: Thus by Corollary 4.5d, the matrix entries $f_{cd}(i)$ satisfy relations (5)-(7).]
c) For $1 \leq c \leq P_i$, define " level j of S_i^c ", or the " j'th level of S_i^c ", to be the $(i-1)$-level $Y_i^{(c-1)2^{e_t - e_{t-1}} + j} \subset Y_i$. Designate level j of S_i^c as being either 'blank' or 'nonblank' as follows.

Level j of S_i^c is 'blank' if $f_{cj}(i) = 1$, and is 'nonblank' if $f_{cj}(i) = 0$.

d) Linearly order all the nonblank $(i-1)$-levels in $S_i^{(P_i)}$ by increasing i'th coordinate; that is, for any two nonblank levels $(i-1)$-levels Y_i^t and $Y_i^{t'}$ we have $Y_i^t < Y_i^{t'}$ if and only if $t < t'$.
e) Let $S_i^c(G)$ (resp. $S_i^{(c)}(G)$) be the set points of $S_i^{(c)}$ lying in nonblank $(i-1)$-levels of S_i^c (resp. $S_i^{(c)}$).

3. [**The Inflation Step**]
a) "Inflate" the image $f_i(G)$ by the map $I_i : Y_i^{(u_i(G))} \to S_i^{(P_i)}(G)$ as follows. Take any $z = (z_1, z_2, \ldots, z_i) \in Y_i^{(u_i(G))}$. Then let

$$I_i(z) = (z_1, z_2, \ldots, z_{i-1}, z_i'),$$

where $Y_i^{z_i'}$ is the z_i'th nonblank $(i-1)$-level in $S_i^{(P_i)}$ (in the ordering of nonblank $(i-1)$-levels in $S_i^{(P_i)}$ from step 2(d)).

[Comment: Let b and r be the integers such that $Y_i^{z_i'}$ is the b'th nonblank $(i-1)$-level of $S_i^r(G)$. Recall that $N_r(b)$ is the column index in matrix $F(i)$ of the b'th zero in row r of $F(i)$. Then by step 2(c), we have $z_i' = (r-1)2^{e_i - e_{i-1}} + N_r(b)$, $1 \le N_r(b) \le 2^{e_i - e_{i-1}}$, so

$$I_i(z) = (z_1, z_2, \ldots, z_{i-1}, (r-1)2^{e_t - e_{t-1}} + N_r(b)).]$$

b) Let $S_i^r(G)' = S_i^r(G) \cap (I_i \circ f_i)(G)$ and $S_i^{(r)}(G)' = S_i^{(r)}(G) \cap (I_i \circ f_i)(G)$.

4. [**The Stacking Step**]

"Stack" the sets $S_i^r(G)'$, $1 \le r \le P_i$, from step 3 (The Inflation Step) on top of each other "over" S_i^1 in succession as r increases. We do this by the stacking map $\sigma_i : S_i^{(P_i)}(G)' \to S_i^1 \times P(u_{i+1}) \subseteq Y_{i+1}^{(u_{i+1})}$ defined as follows. Let $y = (y_1, y_2, \ldots, y_{i-1}, y_i) \in S_i^{(P_i)}(G)'$, say with $y \in S_i^r(G)'$, $1 \le r \le P_i$.

a) Let $n_y = |\{z \in S_t^{(r)}(G)' : z_{1 \to i-1} = y_{1 \to i-1} \text{ and } z_i \equiv y_i \pmod{2^{e_i - e_{i-1}}}\}|$.

b) Let $\bar{y}_i$ be such that $\bar{y}_i \equiv y_i \pmod{2^{e_i - e_{i-1}}}$, with $1 \le \bar{y}_i \le 2^{e_i - e_{i-1}}$. So $\bar{y}_i = N_r(b)$ as in the comment to step 3 (The Inflation Step).

c) Define $\sigma_i(y) = (y_1, y_2, \ldots, y_{i-1}, \bar{y}_i, n_y) = (y_1, y_2, \ldots, y_{i-1}, N_r(b), n_y)$.

[Comment: Since $\sigma_i(y)_{1 \to i} = (y_1, y_2, \ldots, y_{i-1}, \bar{y}_i) \in S_i^1$, we see that $\sigma_i(S_i^{(P_i)}(G)') \subset S_i^1 \times P(m)$, where m is the maximum of n_y over all y belonging to the last set $S_i^{P_i}(G)'$. We will see later that $m = u_{i+1}(G)$.]

5. [**The Composition Step**]

Finally define $f_{i+1} : G \to Y_{i+1}^{(u_{i+1})}$ as the composition $f_{i+1} = \sigma_i \circ I_i \circ f_i$. In particular, for any $z = f_i(v) \in f_i(G)$ we have $f_{i+1}(v) = \sigma_i(I_i(z))$.

Consider now the construction of f_i from f_{i-1} by the above construction. The stacking step 4 suggests that we can regard $f_i(G) \subseteq S_{i-1}^1 \times P(m)$ (as in the comment to step 4) as a collection of stacks addressed by the points of $S_{i-1}^1 \cong \langle Y_{i-1} \rangle$. The height of each such stack extends into the i'th dimension of Y_i. Specifically, for any $x \in \langle Y_{i-1} \rangle$, we let $Stack_i(x, r) = \{\sigma_{i-1}(y) : (\sigma_{i-1}(y))_{1 \to i-1} = x \text{ and } y \in S_{i-1}^{(r)}(G)'\}$, the stack addressed by x, for a given integer $r \le P_{i-1}$. So $Stack_i(x, r)$ consists of images $\sigma_{i-1}(y)$ which project onto x in their first $i-1$ coordinates, and such that y comes from the first r many sets $S_{i-1}^j(G)'$, $1 \le j \le r$. These sets $S_{i-1}^j(G)'$, $1 \le j \le r$, are stacked on top of $S_{i-1}^1 \cong \langle Y_{i-1} \rangle$ in order of increasing j. We view the "height" of $\sigma_{i-1}(y)$ in $Stack_i(x, r)$ as its i'th coordinate $\sigma_{i-1}(y)_i = n_y$, and the height of $Stack_i(x, r)$ as $|Stack_i(x, r)|$. Now define $[r]_i = \max\{|Stack_i(x, r)| : x \in \langle Y_{i-1} \rangle\}$, the maximum height of any of these stacks addressed by points of $\langle Y_{i-1} \rangle$. Thus by definition $\sigma_{i-1}(S_{i-1}^{(r)}(G)')$ is contained in the first $[r]_i$ many $(i-1)$-levels of Y_i; that is, $\sigma_{i-1}(S_{i-1}^{(r)}(G)') \subset Y_i^{([r]_i)}$. As a convenience, for a stack address $x \in S_{i-1}^1 \cong \langle Y_{i-1} \rangle$, we will refer to x either as a member of S_{i-1}^1 or of $\langle Y_{i-1} \rangle$, in most cases of $\langle Y_{i-1} \rangle$.

The parameter $[r]_i$ plays a role in our containment result. We will see in Lemma 5.1f that $I_{i-1}(f_{i-1}(D_{i-1}^{(r)})) \subseteq S_{i-1}^{(r)}(G)$, as suggested in section 3. Thus $[r]_i \ge \lceil \frac{|D_{i-1}^{(r)}|}{|\langle Y_{i-1} \rangle|} \rceil = \lceil \frac{|D_{i-1}^{(r)}|}{2^{e_{i-1}}} \rceil$. We will also see (Lemma 5.1e1) that in fact equality holds in the first inequality for each relevant r and i; that is, $[r]_i$ is as small as it could possibly be. Thus taking $r = P_{i-1}$ and recalling that $D_{i-1}^{(P_{i-1})} = G$, we obtain $f_i(G) \subseteq Y_i^{(u_i(G))} \subseteq Opt(G)$ for each $i \ge 2$, our containment result.

The stacking step 4 of the construction will yield the following monotonicity properties of $Stack_i(x,r)$. First, if $w', w'' \in Stack_i(x,r)$ with $\sigma_{i-1}^{-1}(w') \in S_{i-1}^c$ and $\sigma_{i-1}^{-1}(w'') \in S_{i-1}^d$, $1 \le c, d \le r$, then $w_i' > w_i''$ implies that $c > d$. That is, the originating $(i-1)$-section number under σ_{i-1}^{-1} is strictly increasing as we move up any fixed stack. For the second property, let $Stack_i'(x,r) = f_i(D_{i-1}^{(r)}) \cap Stack_i(x,r)$. We will see that $Stack_i'(x,r)$ is an initial substack of $Stack_i(x,r)$. Specifically, if $w', w'' \in Stack_i'(x,r)$ with $f_i^{-1}(w') \in D_{i-1}^c$ and $f_i^{-1}(w'') \in D_{i-1}^d$, then $w_i' > w_i''$ implies that $c \ge d$. Thus the originating $(i-1)$-page number under f_i^{-1} is nondecreasing as we move up any fixed stack. The first monotonicity property (which we call *stack monotonicity*) is immediate from our construction, and will be noted for the record in the Lemma which follows (part g). The second monotonicity property (which we call *page monotonicity*) will be proved later in Theorem 6.1(a).

Lemma 5.1 (a) *For $x \in V(G)$. For $2 \le j \le k$ the maps f_j satisfy the following.*
(a1) For $1 \le i \le k-1$ and $i+1 \le j \le k$ we have $f_j(x)_{1\to i} = f_{i+1}(x)_{1\to i}$.
(a2) In particular, suppose $2 \le i \le k-1$ and take $z = f_i(x)$. Let $I_i(z) = (z_1, z_2, \ldots, z_i')$ as in step 3 (the Inflation Step) of the above construction, and express z_i' as $z_i' = (r-1)2^{e_i - e_{i-1}} + N_r(b)$ as in the comment to step 3.
Then $f_k(x)_i = N_r(b)$, and $1 \le f_k(x)_i \le 2^{e_i - e_{i-1}}$.

(b) *For any $x, y \in V(G)$, if $|f_i(x)_i - f_i(y)_i| \le e$ then $|f_k(x)_i - f_k(y)_i| \le 2e+2$, where the last difference is interpreted mod $2^{e_i - e_{i-1}}$.*

(c) *Consider the map $\bar\sigma_i : y \to \sigma_i(y)_{1\to i}$, with $y \in S_i^{(P_i)}(G)'$, obtained from σ_i by projecting onto the first i coordinates. Then $\bar\sigma_i$ is one to one when restricted to any one i-section; that is, to the set $\{y \in S_i^r(G)'\}$ for a given $1 \le r \le P_i$. Hence σ_i is one to one, and f_i is one to one for all $2 \le i \le k$.*

(d) *Let $1 \le r < P_i$ and $1 \le t < P_{i-1}$ be integers. Then*
(d1) For $i \ge 2$, $S_i^{(r)}(G)' = S_i^{(r)}(G)$
(d2) For $i \ge 3$, $|Stack_i(x,t)| = [t]_i$ or $[t]_i - 1$ for any stack address $x \in \langle Y_{i-1}\rangle$.
(d3) For $i \ge 3$ and any two stack addresses $x, y \in \langle Y_{i-1}\rangle$, if $x_{i-1} = y_{i-1}$, then $|Stack_i(x,t)| = |Stack_i(y,t)|$.

(e) *Let $l(i,r) = \lceil \frac{|D_{i-1}^{(r)}|}{2^{e_{i-1}}} \rceil = \lceil \frac{r a_{i-1} a_{i-2} \cdots a_1}{2^{e_{i-1}}} \rceil$, and $l'(i,r) = \lceil \frac{|D_i^{(r)}|}{2^{e_{i-1}}} \rceil = \lceil \frac{r a_i a_{i-1} \cdots a_1}{2^{e_{i-1}}} \rceil$. Then for $i \ge 3$*
(e1) for $1 \le r \le P_{i-1}$ we have $[r]_i = l(i,r)$, and $|Y_i^{(l(i,r))}| - |f_i(D_{i-1}^{(r)})| < 2^{e_{i-1}}$, and
(e2) for $1 \le r \le P_{i-1}$ we have $f_i(D_{i-1}^{(r)}) \subseteq Y_i^{(l(i,r))}$, and for $1 \le r \le P_i$, $f_i(D_i^{(r)}) \subseteq Y_i^{(l'(i,r))}$.
(e3) $f_i(G) \subseteq Y_i^{(u_i)} \subseteq Opt(G)$. Moreover, we have $f_k(G) \subseteq Opt'(G)$ and $H^k(G) \subseteq Opt(G)$.

(f) *For $i \ge 2$ we have $I_i(f_i(D_i^{(r)})) \subseteq S_i^{(r)}(G)$ for $1 \le r \le P_i$, and $S_i^{(r)}(G) \subset I_i(f_i(D_i^{(r+1)}))$ for $1 \le r \le P_i - 1$.*

(g) *(Stack monotonicity) Suppose $w', w'' \in Stack_i(x,r)$, $1 \le r \le P_{i-1}$, with $\sigma_{i-1}^{-1}(w') \in S_{i-1}^c$ and $\sigma_{i-1}^{-1}(w'') \in S_{i-1}^d$, $1 \le c, d \le r$. Then $w_i'' > w_i' \Leftrightarrow d > c$.*

Proof. Consider first part (a1). We proceed by induction on j. The base case $j = i+1$ is trivial. So suppose inductively that $f_j(x)_{1\to i} = f_{i+1}(x)_{1\to i}$ for some $j \ge i+1$. By the definition of I_j and σ_j from the inflation and stacking steps respectively, we have $I_j(z)_{1\to j-1} = z_{1\to j-1}$ and $\sigma_j(y)_{1\to j-1} = y_{1\to j-1}$ for z and y in the domain of I_j and σ_j respectively. Thus since $j \ge i+1$

we have $I_j(f_j(x))_{1\to i} = f_j(x)_{1\to i}$. Therefore $f_{j+1}(x)_{1\to i} = \sigma_j(I_j(f_j(x)))_{1\to i} = I_j(f_j(x))_{1\to i} = f_j(x)_{1\to i} = f_{i+1}(x)_{1\to i}$. This completes the inductive step, so (a1) is proved.

Next consider (a2). For $i \geq 2$ we have $f_{i+1}(x)_i = N_r(b)$ by step 4c (The Stacking Step), and $1 \leq N_r(b) \leq 2^{e_i - e_{i-1}}$ by the comment to step 3 and the definition of $N_r(b)$, proving (a2).

Consider (b). We know that $I_i(f_i(x))$ is in some nonblank $(i-1)$-level, say the b'th one, of some i-section, say S_i^r, of $S_i^{(P_i)}$. So we can write $I_i(f_i(x))_i = (r-1)2^{e_i-e_{i-1}} + N_r(b)$. Since $|f_i(x)_i - f_i(y)_i| \leq e$ and I_i preserves the order of $(i-1)$-levels, it follows that $|I_i(f_i(x))_i - I_i(f_i(y))_i| \leq e$. Assuming without loss that $I_i(f_i(y))_i > I_i(f_i(x))_i$ (otherwise interchange the roles of x and y), we can write $I_i(f_i(y))_i = (r-1)2^{e_i-e_{i-1}} + N_r(b+t)$ for some $0 \leq t \leq e$, where we interpret the function $N_r(*)$ in row major order as in the discussion just preceding Corollary 4.5. So we have $|f_k(x)_i - f_k(y)_i| \leq |N_r(b+e) - N_r(b)| \leq 2e+2$, where the first inequality follows from part (a2), and the second by using the more generous of the bounds (a) and (b) in Corollary 4.5, and interpreting the differences mod $2^{e_i-e_{i-1}}$.

Consider (c). For the first statement, let $y, z \in S_i^r(G)'$ for some $1 \leq r \leq P_i$. We must show that $(\sigma_i(y))_{1\to i} \neq (\sigma_i(z))_{1\to i}$. By the comment to step 3 (The Inflation Step) we may write $y = (y_1, y_2, \ldots, y_{i-1}, (r-1)2^{e_i-e_{i-1}} + N_r(d))$ and $z = (z_1, z_2, \ldots, z_{i-1}, (r-1)2^{e_i-e_{i-1}} + N_r(c))$ for suitable integers d and c and $1 \leq N_r(d), N_r(c) \leq 2^{e_i-e_{i-1}}$. By step 4 (The Stacking Step), we have $(\sigma_i(z))_{1\to i} = (z_1, z_2, \ldots, z_{i-1}, N_r(d))$, and $(\sigma_i(y))_{1\to i} = (y_1, y_2, \ldots, y_{i-1}, N_r(c))$. If y and z disagree at one of their first $i-1$ coordinates, then obviously $(\sigma_i(y))_{1\to i} \neq (\sigma_i(z))_{1\to i}$ by these formulas. So we can suppose that $y_{1\to i-1} = z_{1\to i-1}$. But since y and z are distinct, the formulas above for y and z force them to disagree in their i'th coordinates, so $N_r(d) \neq N_r(c)$. Thus $y_i \not\equiv z_i \pmod{2^{e_i-e_{i-1}}}$. So by definition of σ_i we get $(\sigma_i(y))_{1\to i} \neq (\sigma_i(z))_{1\to i}$, as desired. For one-to-oneness of σ_i itself, it only remains to show that if y, z come from distinct sections, then $\sigma_i(y) \neq \sigma_i(z)$. So let $y \in S_i^s(G)'$ and $z \in S_i^t(G)'$, say with $s < t$. If $(\sigma_i(y))_{1\to i} \neq (\sigma_i(z))_{1\to i}$, then the claim follows obviously, so assume $(\sigma_i(y))_{1\to i} = (\sigma_i(z))_{1\to i}$. Then $\sigma_i(y)$ and $\sigma_i(z)$ both belong to the stack addressed by $(\sigma_i(y))_{1\to i}$, namely, $Stack_{i+1}((\sigma_i(y))_{1\to i}, t)$. But then since $s < t$, $\sigma_i(y)$ is lower in this stack than $\sigma_i(z)$ by definition of σ_i. Then $(\sigma_i(y))_{i+1} < (\sigma_i(z))_{i+1}$, proving one-to-oneness of σ_i. As for the claim about the f_i, observe first that f_2 is one-to-one. So from $f_{i+1} = \sigma_i \circ I_i \circ f_i$ and the one-to-oneness of σ_i and I_i, it follows by induction on i that f_i is one-to-one for all $2 \leq i \leq k$.

For (d), we start by proving (d1) for $i = 2$. By Theorem 2.2e,f, $Y_2^{(m-1)} \subset f_2(G) \subseteq Y_2^m$, where $m = \lceil \frac{|G|}{2^{e_1}} \rceil$ (noting that $m = u_2(G)$). That is, all but at most the last column Y_2^m of $Y_2^{(m)}$ is contained in $f_2(G)$. Applying I_2 to both sides of the last containment and noting that $(I_2 \circ f_2)(G) = S_2^{(P_2)}(G)'$, we see that every nonblank column of $S_2^{(P_2)}(G)$ lies in $S_2^{(P_2)}(G)'$, except possibly the last one $I_2(Y_2^m)$, call it Y_2^d. Since every 2-section must contain at least one nonblank column we have $Y_2^d \subset S_2^{P_2}$ (the last 2-section), so $S_2^{(P_2-1)}(G) = S_2^{(P_2-1)}(G)'$. Taking subsets, we get $S_2^{(r)}(G) = S_2^{(r)}(G)'$ for $1 \leq r < P_2$.

Next we show that (d1) for i implies (d2) and (d3) for $i+1$. Take $x = (x_1, x_2, \ldots, x_i) \in \langle Y_i \rangle$ to be a stack address. By definition of matrix $F(i)$, its entry $f_{t,x_i}(i)$ satisfies $f_{t,x_i}(i) = 0$ if and only if the $(i-1)$-level $Y = Y_i^{(t-1)2^{e_i-e_{i-1}}+x_i}$ (the x_i'th $(i-1)$-level of i-section S_i^t) is nonblank. Assume first that $f_{t,x_i}(i) = 0$. Thus Y is nonblank. Since $t < P_i$ (by the hypothesis for (d2) with $i+1$ replacing i), we have $Y \subset S_i^{(P_i-1)}(G) = S_i^{(P_i-1)}(G)'$, the last equality holding by the conclusion of (d1) for i. Thus every point of Y lies in the domain of

σ_i. So $Stack_{i+1}(x,t)$ receives the point $\sigma_i(x_1, x_2, \ldots, (t-1)2^{e_i - e_{i-1}} + x_i)$ under the stacking map σ_i. If $f_{t,x_i}(i) = 1$, then $Stack_{i+1}(x,t)$ receives no point $\sigma_i(y)$ with $y \in S_i^t$. It follows that $|Stack_{i+1}(x,t)| = t - \sum_{j=1}^{t} f_{j,x_i}(i)$. Since by Lemma 4.4b the sum on the right must be one of two successive integers depending on t, it follows that $|Stack_{i+1}(x,t)|$ is one of two successive integers depending on t but independent of x. Thus by definition of $[t]_{i+1}$, we get $|Stack_{i+1}(x,t)| = [t]_{i+1}$ or $[t]_{i+1} - 1$, proving (d2) for $i+1$. Consider now (d3) for $i+1$, and let $y \in \langle Y_i \rangle$ be a stack address with $y_i = x_i$. Then clearly $f_{j,y_i}(i) = f_{j,x_i}(i)$ for $1 \le j \le P_i$ by the definition of matrix $F(i)$. Thus $\sum_{j=1}^{t} f_{j,x_i}(i) = \sum_{j=1}^{t} f_{j,x_i}(i)$, and hence $|Stack_{i+1}(x,t)| = |Stack_{i+1}(y,t)|$, as required.

Finally we show that (d2) for $i+1$ in place of i implies (d1) for $i+1$ in place of i. By assumption we have $|Stack_{i+1}(x,t)| = [t]_{i+1}$ or $[t]_{i+1} - 1$ for $t \le P_i - 1$ and any stack address $x \in \langle Y_i \rangle$. Observe that $[P_i]_{i+1} \le [P_i - 1]_{i+1} + 1$ since the projection σ_i onto the first i coordinates is one-to-one when restricted to any single section, in this case $S_i^{P_i}$, by part (c). Further, $|Stack_{i+1}(x, P_i - 1)| \ge [P_i - 1]_{i+1} - 1$ for all stack addresses x by assumption. It follows that $Y_{i+1}^{([P_i - 1]_{i+1} - 1)} \subseteq f_{i+1}(G) \subseteq Y_{i+1}^{([P_i]_{i+1})} \subseteq Y_{i+1}^{([P_i - 1]_{i+1} + 1)}$. So at most two i-levels of $Y_{i+1}^{([P_i]_{i+1})}$ are possibly not entirely contained in $f_{i+1}(G)$, these being the top two $Y_{i+1}^{[P_i]_{i+1}}$ and $Y_{i+1}^{[P_i]_{i+1} - 1}$, and if this possibility occurs then $[P_i]_{i+1} = [P_i - 1]_{i+1} + 1$ (so also $[P_i]_{i+1} - 1 = [P_i - 1]_{i+1}$). But still $Y_{i+1}^{j} \cap f_{i+1}(G) \ne \emptyset$ if and only if $1 \le j \le [P_i]_{i+1}$ by definition of $[P_i]_{i+1}$. Thus every i-level $I_{i+1}(Y_{i+1}^{j})$, $1 \le j \le [P_i]_{i+1}$, is nonblank, and by the previous sentence all but at most two of these nonblank i-levels belong to $S_{i+1}^{(P_{i+1})}(G)'$, the two possible exceptions being $I_{i+1}(Y_{i+1}^{[P_i]_{i+1}})$ and $I_{i+1}(Y_{i+1}^{[P_i]_{i+1} - 1})$. Since any $(i+1)$-section S_{i+1}^{j}, $1 \le j \le P_{i+1}$, contains at least two nonblank i-levels (since $a_i > 4$ for all i) it follows that the top two nonblank i-levels of $S_{i+1}^{(P_{i+1})}$ belong to the top $(i+1)$-section $S_{i+1}^{P_{i+1}}$. So the remaining $(i+1)$-sections S_{i+1}^{j}, $j < P_{i+1}$, satisfy $S_{i+1}^{j}(G) = S_{i+1}^{j}(G)'$. In particular we have $S_{i+1}^{(P_{i+1} - 1)}(G) = S_{i+1}^{(P_{i+1} - 1)}(G)'$, and taking subsets we obtain (d1) for $i+1$.

Consider now (e1). The case $i = 2$ follows immediately from Theorem 2.2e,f, and we assume $i \ge 3$.

For the lower bound $[r]_i \ge \lceil \frac{|D_{i-1}^{(r)}|}{2^{e_i - 1}} \rceil = l(i,r)$, suppose first that $r < P_{i-1}$. By definition we have $[r]_i \ge \frac{|\sigma_i(S_{i-1}^{(r)}(G)')|}{|\langle Y_{i-1} \rangle|} = \frac{|\sigma_i(S_{i-1}^{(r)})(G)')|}{2^{e_i - 1}}$. So since σ_i is one to one, it suffices to show that $|S_{i-1}^{(r)}(G)'| \ge \lceil \frac{|D_{i-1}^{(r)}|}{2^{e_i - 2}} \rceil 2^{e_i - 2}$. For then we get $[r]_i \ge \lceil \frac{|D_{i-1}^{(r)}|}{2^{e_i - 2}} \rceil 2^{e_i - 2 - e_{i-1}} \ge \frac{|D_{i-1}^{(r)}|}{2^{e_i - 1}}$, and since $[r]_i$ is an integer $[r]_i \ge \lceil \frac{|D_{i-1}^{(r)}|}{2^{e_i - 1}} \rceil = l(i,r)$. By part (d1) and Lemma 4.1 we have $S_{i-1}^{(r)}(G) = S_{i-1}^{(r)}(G)'$, and $S_{i-1}^{(r)}(G)'$ consists of $\lceil \frac{|D_{i-1}^{(r)}|}{2^{e_i - 2}} \rceil$ many $(i-2)$-levels in Y_{i-1}. Since each such $(i-2)$-level has $2^{e_i - 2}$ points, we get $|S_{i-1}^{(r)}(G)'| \ge \lceil \frac{|D_{i-1}^{(r)}|}{2^{e_i - 2}} \rceil 2^{e_i - 2}$ as required. Now suppose that $r = P_{i-1}$. We have $|S_{i-1}^{(P_{i-1})}(G)'| = |G| = |P_{i-1} a_{i-1} a_{i-2} \cdots a_1|$. So again since σ_i is one to one we get $[P_{i-1}]_i \ge \frac{|\sigma_i(S_{i-1}^{(P_{i-1})}(G)')|}{|\langle Y_{i-1} \rangle|} \ge \frac{|P_{i-1} a_{i-1} a_{i-2} \cdots a_1|}{2^{e_i - 1}} = \frac{|D_{i-1}^{(P_{i-1})}|}{2^{e_i - 1}}$, as required.

It remains to show that $[r]_i \le l(i,r)$. Recall that for any stack address $x \in \langle Y_{i-1} \rangle$, $|Stack_i(x,r)|$ is at most the number of 0's in column x_i among the first r rows in matrix $F(i-1)$. By Theorem 4.4b this number of 0's is either the same for all x or is one of two successive integers, call them α_r or $\alpha_r - 1$, depending only on r (and i, which we fix in this argument). Since $[r]_i$ is the maximum of $|Stack_i(x,r)|$ over all $x \in \langle Y_{i-1} \rangle$, it suffices to prove

that $\alpha_r \le l(i,r)$. By our construction, the total number of 0's in the first r rows of $F(i-1)$ is the number of of nonblank $(i-2)$-levels in $S_{i-1}^{(r)}$. So by Lemma 4.1a this number of 0's is $\lceil \frac{ra_{i-1}a_{i-2}\cdots a_1}{2^{e_{i-2}}} \rceil$. Let $\frac{p}{2^{e_{i-2}}} = \lceil \frac{ra_{i-1}a_{i-2}\cdots a_1}{2^{e_{i-2}}} \rceil - \frac{ra_{i-1}a_{i-2}\cdots a_1}{2^{e_{i-2}}}$. Since the number of columns of $F(i-1)$ is $2^{e_{i-1}-e_{i-2}}$, we have $\alpha_r = \lceil \frac{\lceil \frac{ra_{i-1}a_{i-2}\cdots a_1}{2^{e_{i-2}}} \rceil}{2^{e_{i-1}-e_{i-2}}} \rceil = \lceil \frac{ra_{i-1}a_{i-2}\cdots a_1}{2^{e_{i-1}}} + \frac{p}{2^{e_{i-1}}} \rceil = \lceil \frac{ra_{i-1}a_{i-2}\cdots a_1}{2^{e_{i-1}}} \rceil$, as required.

For the second claim of (e1), again note that each $(i-1)$-level of Y_i has size $2^{e_{i-1}}$. Thus $|Y_i^{(l(i,r))}| - |f_i(D_{i-1}^{(r)})| = 2^{e_{i-1}} \lceil \frac{|D_{i-1}^{(r)}|}{2^{e_{i-1}}} \rceil - |D_{i-1}^{(r)}| < 2^{e_{i-1}}$, completing the proof of (e1).

For (e2) we prove both $f_i(D_{i-1}^{(r)}) \subseteq Y_i^{(l(i,r))}$ and $f_i(D_i^{(r)}) \subseteq Y_i^{(l'(i,r))}$ by induction. If $i = 2$, then both statements hold by Theorem 2.2d,e,f. So let $i > 2$ be given, and assume inductively that both of these containments hold for $i-1$ in place of i. By this assumption we have $f_{i-1}(D_{i-1}^{(r)}) \subseteq Y_{i-1}^{(l'(i-1,r))}$. Applying the inflation map I_{i-1} to both sides, we obtain $(I_{i-1} \circ f_{i-1})(D_{i-1}^{(r)}) \subseteq S_{i-1}^{(r)}(G)$, since by Lemma 4.1 we see that $S_{i-1}^{(r)}$ has the same number $l'(i-1,r)$ of nonblank $(i-2)$-levels as the number of $(i-2)$-levels in $Y_{i-1}^{(l'(i-1,r))}$, and I_{i-1} preserves the order of the latter set of $(i-2)$-levels. Thus $(I_{i-1} \circ f_{i-1})(D_{i-1}^{(r)}) \subseteq S_{i-1}^{(r)}(G)'$ by definition of $S_{i-1}^{(r)}(G)'$. Now applying σ_{i-1} to the last containment, and recalling that $f_i = \sigma_{i-1} \circ I_{i-1} \circ f_{i-1}$, we get $f_i(D_{i-1}^{(r)}) = (\sigma_{i-1} \circ I_{i-1})(f_{i-1}(D_{i-1}^{(r)})) \subseteq \sigma_{i-1}(S_{i-1}^{(r)}(G)') \subseteq Y_i^{([r]_i)}$, where the last containment holds by definition of $[r]_i$. By (e1) we have $[r]_i = l(i,r)$, thereby proving $f_i(D_{i-1}^{(r)}) \subseteq Y_i^{(l(i,r))}$. To show $f_i(D_i^{(r)}) \subseteq Y_i^{(l'(i,r))}$, apply $f_i(D_{i-1}^{(r)}) \subseteq Y_i^{(l(i,r))}$ using $a_i r$ in place of r and observing that $D_i^{(r)} = D_{i-1}^{(a_i r)}$. This completes the inductive step, and hence the proof of (e2).

Consider now (e3). Recall that $P_{i-1} = \prod_{t=i}^{k} a_t$, so that $V(G) = V(D_{i-1}^{(P_{i-1})})$. Noting that $l(i, P_{i-1}) = \lceil \frac{|D_{i-1}^{(P_{i-1})}|}{2^{e_{i-1}}} \rceil = \lceil \frac{|V(G)|}{2^{e_{i-1}}} \rceil = u_i(G)$, by part (e2) we have $f_i(G) \subseteq Y_i^{(l(i,P_{i-1}))} = Y_i^{(u_i(G))}$, proving the first containment. For the rest, note that $u_i(G) \le 2^{\lceil log_2(|G|) \rceil - e_{i-1}}$, so $f_i(G) \subseteq Y_i^{(u_i(G))} \subseteq \langle Y_{i-1} \rangle \times P(2^{\lceil log_2(|G|) \rceil - e_{i-1}})$. The last graph is a spanning subgraph of $Opt(G)$ for each i, yielding the second containment, and for $i = k$ yielding the final two containments after performing the relabeling with hypercube addresses by the map H^k.

For (f), start with $f_i(D_i^{(r)}) \subseteq Y_i^{(l'(i,r))}$ from (e2). Then we get $I_i(f_i(D_i^{(r)})) \subseteq I_i(Y_i^{(l'(i,r))}) = S_i^{(r)}(G)$. The equality holds since the right side consists of $l'(i,r)$ many (nonblank) $(i-1)$-levels, while I_i preserves the order of $(i-1)$-levels by increasing i-coordinate value.

Next we show that $S_i^{(r)}(G) \subset I_i(f_i(D_i^{(r+1)}))$. For the case $i = 2$ we apply Theorem 2.2e,f and Corollary 2.3e to $G = D_2^{(r+1)}$, together with Lemma 4.1 to obtain $S_2^{(r)}(G) = I_2(Y_2^{(l'(2,r))}) \subseteq I_2(Y_2^{(l'(2,r+1)-1)}) \subseteq I_2(f_2(D_2^{(r+1)}))$.

Proceeding by induction on i, let $i \ge 3$ and assume the statement true for $i-1$ and $1 \le r \le P_{i-1} - 1$, and we prove it for i and $1 \le r \le P_i - 1$. It suffices to show that $Y_i^{(l'(i,r))} \subset f_i(D_i^{(r+1)})$, since then we can just apply I_i to each side and use $I_i(Y_i^{(l'(i,r))}) = S_i^{(r)}(G)$ (as observed above) to get the result directly.

By (e1) we have $[a_i r + 2]_i = l(i, a_i r + 2) \ge l'(i,r) + 1$, using $2a_{i-1}a_{i-2} \ldots a_1 > 2^{e_{i-1}}$. By (d), for any $x \in \langle Y_{i-1} \rangle$ we have $|Stack_i(x, a_i r + 2)| = [a_i r + 2]_i$ or $[a_i r + 2]_i - 1$, so $|Stack_i(x, a_i r + 2)| \ge l'(i,r)$. It follows that $Y_i^{(l'(i,r))} \subseteq \sigma_{i-1}(S_{i-1}^{(a_i r+2)}(G)') \subseteq \sigma_{i-1}(I_{i-1}(f_{i-1}(D_{i-1}^{(a_i r+3)})))$, where the last containment is by the inductive hypothesis. Now since $a_i \ge 2^{22} > 3$, we have $a_i r + 3 < a_i(r+1)$, so $D_{i-1}^{(a_i r+3)} \subset D_{i-1}^{(a_i(r+1))} = D_i^{(r+1)}$. So we have $Y_i^{(l'(i,r))}(G) \subset (\sigma_{i-1} \circ I_{i-1} \circ f_{i-1})(D_i^{(r+1)}) =$

$f_i(D_i^{(r+1)})$, as desired.

For (g) observe that σ_{i-1} stacks the sets $S_{i-1}^r(G)$ onto S_{i-1}^1 in succession by increasing r; that is, $S_{i-1}^c(G)$ is stacked before $S_{i-1}^d(G)$ precisely when $c < d$. Part (g) follows. ■

We can now give the final embedding H^k of G into $Opt'(G)$. It is based on the fact that $f_k(G) \subseteq Opt'(G)$ from Lemma 5.1(e3). So by definition we have $1 \leq f_k(x)_j \leq 2^{e_j - e_{j-1}}$ for $1 \leq j \leq k$ and all $x \in V(G)$. Given these facts, we obtain H^k from f_k as follows. For each $1 \leq j \leq k$ we interpret $f_k(x)_j$ as a hypercube point in $Q_{e_j - e_{j-1}}$ using the inverse image of the labeling $L_{e_j - e_{j-1}}$ of Corollary 4.8. We then concatenate these hypercube points (now strings over $\{0,1\}$) left to right in order of increasing j to obtain $H^k(x)$. The details are as follows.

Construction of the map $H^k : G = [a_1 \times a_2 \times a_3 \times \cdots \times a_k] \to Opt(G)$

1. Initialization and the case $k = 2$.
a) Start with the map $f_2 : G \to Y_2^{u_2} \subseteq P(2^{e_1}) \times P(2^{e_2 - e_1})$.
b) Define $H^2 : G \to Opt(G)$ by $H^2(x) = (L_{e_1}^{-1}(f_2(x)_1), L_{e_2-e_1}^{-1}(f_2(x)_2))$.
2. For $k \geq 3$, construct the maps $f_3, f_4, \ldots, f_k$ inductively as follows.
For $i = 3$ to $k-1$, construct $f_{i+1} : G \to Y_{i+1}$ from f_i using the procedure given at the beginning of this subsection.
3. Having obtained the map $f_k : G \to Opt'(G)$ from the preceding step, define the map $H^k : G \to Opt(G)$ by

$H^k(x) = (H^k(x)_1, H^k(x)_2, \ldots, H^k(x)_k)$, where $H^k(x)_j = L_{e_j - e_{j-1}}^{-1}(f_k(x)_j)$ for $1 \leq j \leq k$,

taking $e_0 = 0$, and where $L_{e_j - e_{j-1}}$ is the labeling from Corollary 4.8.

6 The dilation bound

From Lemma 5.1e3 we have the containment result $H^k(G) \subseteq Opt(G)$. The goal in this section is to complete the proof of our main result by showing that $dilation(H^k) \leq 3k$ when every a_i is larger than some fixed constant.

We recall some notation. For $x \in \langle Y_{i-1} \rangle$, recall that $Stack_i(x,r) = \{z = \sigma_{i-1}(y) : z_{1 \to i-1} = x \text{ and } y \in S_{i-1}^{(r)}(G)\}$, that $Stack'_i(x,r) = f_i(D_{i-1}^{(r)}) \cap Stack_i(x,r)$, and that $[r]_i = \max\{|Stack_i(x,r)| : x \in \langle Y_{i-1} \rangle\}$. So by definition we have $Stack'_i(x,r) \subseteq Stack_i(x,r)$.

To set the context for the next theorem, note that by Lemma 5.1e1,e2, $f_i(D_{i-1}^{(r)}) \subseteq Y_i^{([r]_i)}$, so that $|Stack'_i(x,r)| \leq [r]_i$ In the next theorem we will see that $Stack'_i(x,r)$ is always an initial substack of $Stack_i(x,r)$; equivalently, that $Stack'_i(x, P_{i-1})$ is page monotone. Also we will see that the "page stack" heights $|Stack'_i(x,r)|$, $x \in \langle Y_{i-1} \rangle$, fall within a narrow range; for a given r (and fixed i) any two such heights differ by at most 2 independent of x.

For a subset $S \subseteq Y_i^{([r]_i)}$, let $v_{i,r}(S) = |S \cap f_i(D_{i-1}^{(r)})|$.

Theorem 6.1 *Let $i \geq 3$ and $1 \leq r \leq P_{i-1}$.*
(a) *(page monotonicity) Take $x \in \langle Y_{i-1} \rangle$. Let $z', z'' \in Stack_i(x, P_{i-1})$ with $f_i^{-1}(z') \in D_{i-1}^s$ and $f_i^{-1}(z') \in D_{i-1}^t$. If $z'_i > z''_i$, then $s \geq t$.*
(b) *For every stack address $x \in \langle Y_{i-1} \rangle$, we have $[r]_i - 2 \leq |Stack'_i(x,r)| \leq [r]_i$. Moreover, all points of $Y_i^{([r]_i)} - f_i(D_{i-1}^{(r)})$ lie in the union $Y_i^{[r]_i} \cup Y_i^{[r]_i - 1}$ of the top two $(i-1)$-levels of $Y_i^{([r]_i)}$.*
(c) $v_{i,r}(Y_i^{[r]_i - 1}) + v_{i,r}(Y_i^{[r]_i}) > 2^{e_{i-1}}$.

Proof. Consider part (a). It suffices to show that for any $x \in \langle Y_{i-1}\rangle$ and $1 \le r \le P_{i-1}$, we have that $Stack_i(x,r)$ is page monotone. We prove this by induction on r, for any fixed $i \ge 3$ and $x \in \langle Y_{i-1}\rangle$. For the base case $r = 1$, the claim is trivial since $Stack_i(x,1)$ contains a single entry by one to oneness of $\bar\sigma_{i-1}$ (Lemma 5.1c) on any one section (in this case, on S^1_{i-1}). So suppose inductively that $Stack_i(x,r)$ is page monotone for some $1 \le r < P_{i-1}$. We use Lemma 5.1f, with $i-1$ in place of i. Applying σ_{i-1} to the second containment $S^{(r)}_{i-1}(G) \subset I_{i-1}(f_{i-1}(D^{(r+1)}_{i-1}))$ stated there and noting that $f_i = \sigma_{i-1} \circ I_{i-1} \circ f_{i-1}$, we see that every entry z of $Stack_i(x,r)$ satisfies $f_i^{-1}(z) \in D^{(r+1)}_{i-1}$. If $Stack_i(x,r+1) = Stack_i(x,r)$, then trivially $Stack_i(x,r+1)$ is page monotone by induction. So assume $Stack_i(x,r+1) \ne Stack_i(x,r)$, and let y be the unique element (by one to oneness of $\bar\sigma_{i-1}$ on any section) of $Stack_i(x,r+1) - Stack_i(x,r)$. By the first containment in Lemma 5.1f (again with $i-1$ replacing i) we have, on applying σ_{i-1} to each side again, $f_i^{-1}(y) \in D^j_{i-1}$ for some $j \ge r+1$. Now $Stack_i(x,r)$ is page monotone by induction and as just noted $f_i^{-1}(z) \in D^{(r+1)}_{i-1}$ for all $z \in Stack_i(x,r)$. Since $Stack_i(x,r+1)$ is obtained by placing y at the top of $Stack_i(x,r)$ and $y \in f_i(D^j_{i-1})$ for some $j \ge r+1$, it follows that $Stack_i(x,r+1)$ is page monotone, completing the inductive step.

For part (b), the upper bound $|Stack'_i(x,r)| \le [r]_i$ follows from $Stack'_i(x,r) \subseteq Stack_i(x,r)$ and the definition of $[r]_i$.

Consider the lower bound on $|Stack'_i(x,r)|$ in part (b). The statement holds vacuously for $r = 1$ and all $i \ge 3$, since $|Stack'_i(x,1)| = 0$ or 1 depending on x, and $[1]_i = 1$, by one to oneness of the projection map $\bar\sigma_{i-1}$ on any section from Lemma 5.1c. So suppose $r \ge 2$.

Fixing i, for any stack address $x \in \langle Y_{i-1}\rangle$, let $s(x,r) = |Stack_i(x,r)|$ and $s'(x,r) = |Stack'_i(x,r)|$. By Lemma 5.1f we have $S^{(r-1)}_{i-1}(G) \subseteq (I_{i-1} \circ f_{i-1})(D^{(r)}_{i-1})$. Now applying σ_{i-1} to both sides of this containment and using $Stack_i(x,r-1) \subseteq Stack_i(x,r)$ and $f_i = \sigma_{i-1} \circ I_{i-1} \circ f_{i-1}$, we have $Stack_i(x,r-1) = [\sigma_{i-1}(S^{(r-1)}_{i-1}(G)) \cap Stack_i(x,r-1)] \subseteq [f_i(D^{(r)}_{i-1}) \cap Stack_i(x,r)] = Stack'_i(x,r)$. Thus $s'(x,r) \ge s(x,r-1)$. Since $r-1 \le P_{i-1}-1$, by Lemma 5.1d we have $s(x,r-1) = [r-1]_i$ or $[r-1]_i - 1$. Therefore $s'(x,r) \ge [r-1]_i - 1$. By Lemma 5.1c we have $[r-1]_i \le [r]_i \le [r-1]_i + 1$. Thus $s'(x,r) \ge [r-1]_i - 1 \ge [r]_i - 2$, proving the first sentence of (b).

The second sentence of (b) follows from the first sentence, together with the page monotonicity property of part (a).

Next consider (c). By Lemma 5.1e1 we have $|Y_i^{([r]_i)}| - |f_i(D^{(r)}_{i-1})| < 2^{e_{i-1}}$. Let $A = |Y_i^{[r]_i} \cup Y_i^{[r]_i - 1}|$ and $B = v_{i,r}(Y_i^{[r]_i - 1}) + v_{i,r}(Y_i^{[r]_i})$. By (b) we have $Y_i^{([r]_i - 2)} \subseteq f_i(D^{(r)}_{i-1})$, so $|Y_i^{([r]_i)}| - |f_i(D^{(r)}_{i-1})| = A - B$. So $2^{e_{i-1}} > |Y_i^{([r]_i)}| - |f_i(D^{(r)}_{i-1})| = A - B = 2 \cdot 2^{e_{i-1}} - B$, so $B > 2^{e_{i-1}}$, as required. ∎

We introduce notation for identifying particular $(i-1)$-subpages of a given i-page in G. For $1 \le j \le a_i$ let $D^r_i(j) = D^{(r-1)a_i + j}_{i-1}$, and we regard $D^r_i(j)$ as the j'th $(i-1)$-subpage of D^r_i under the ordering of $(i-1)$-subpages of D^r_i induced by $\prec_{i-1}$. Similarly let $S^r_i(j)$, $j \ge 1$, be the j'th nonblank $(i-1)$-level of $S^r_i(G)$, ordered by increasing i-coordinate.

Now suppose $z = (I_i \circ f_i)(x)$ for some $x \in D^r_i$. We let $\nu_i(z)$ be the integer such that $z \in S^r_i(\nu_i(z))$ for suitable r; that is, z belongs to the $\nu_i(z)$'th nonblank $(i-1)$-level, ordered by increasing i-coordinate, of the i-section S^r_i which contains z. To illustrate, recall the examples at the right of Figure 5 and the left of Figure 7. In Figure 5 we have the first 8 many 2-levels of Y_3 containing the image $f_3(G)$, where $G = [3 \times 7 \times 4 \times 3]$. In Figure 7 at left we have inserted 4

blank 2-levels among these (as specified by the matrix in Table 1c), and grouped the resulting 12 many 2-levels into the three 3-sections S_3^j, $1 \le j \le 3$, preserving the order of the nonblank levels. So for example the 6'th 2-level in Figure 5 (ordered by 3'rd coordinate, or height in the figure) becomes, after the map I_2 inserts these 4 blank 2-levels, the 3'rd nonblank 2-level $S_3^2(3)$ of section S_3^2 (again ordered by 3'rd coordinate, or height within S_3^2) in Figure 7. So any point $z = (I_3 \circ f_3)(x)$, where $f_3(x)$ was in the 6'th 2-level of Figure 5, satisfies $\nu_3(z) = 3$ since z belongs to the third nonblank 2-level of the 3-section (in this case S_3^2) containing z (as shown in Figure 7). Similarly take any point $f_3(x)$ lying in the 7'th level of Figure 5. Then the corresponding point $z = (I_3 \circ f_3)(x)$ after inflation satisfies $\nu_3(z) = 1$, since z belongs to the first nonblank 2-level $S_3^3(1)$ of the 3-section S_3^3 containing z (as shown in Figure 7).

Let m_r be the number of nonblank $(i-1)$-levels in S_i^r for any $1 \le r \le P_i$. We interpret the $(i-1)$-levels $S_i^r(j)$ for integers j outside the range $[1, m_r]$, and also interpret the differences $\nu_i(z') - \nu_i(z'')$ using "wraparound" as follows. For example, if $j \le 0$, then $S_i^r(j)$ is understood as $S_i^{r-1}(m_{r-1} + j)$. Similarly, if $j > m_r$, then $S_i^r(j)$ is understood as $S_i^{r+1}(j - m_{r+1})$. Also let $z' \in S_i^r(\nu_i(z'))$ and $z'' \in S_i^s(\nu_i(z''))$, where $1 \le \nu_i(z') \le m_r$ and $1 \le \nu_i(z'') \le m_s$. Then define $||\nu_i(z'') - \nu_i(z')||$ as the minimum of $\{|\nu_i(z'') - \nu_i(z')|, m_r - \nu_i(z') + \nu_i(z''), m_s - \nu_i(z'') + \nu_i(z')\}$.

Corollary 6.2 *For $2 \le i \le k$ we have the following.*

(a) *For each $i \ge 2$ and $j \ge 1$, we have $(I_i \circ f_i)(D_i^j) \subseteq S_i^{j-1} \cup S_i^j$. Hence for $1 \le r \le P_{i-1}$ we have the following.*

(1) For any stack address $x \in \langle Y_{i-1} \rangle$ and $1 \le r \le P_{i-1}$, $|Stack_i(x, r) \cap f_i(D_{i-1}^r)| \le 2$.

(2) Suppose $|Stack_i(x, r) \cap f_i(D_{i-1}^r)| = 2$, and let $Stack_i(x, r) \cap f_i(D_{i-1}^r) = \{z', z''\}$. Then z' and z'' are at successive heights in $Stack_i(x, r)$; that is, $|z_i' - z_i''| \le 1$.

(3) Let $z \in D_{i-1}^r$, with $f_i(z) \in Stack_i(x, r)$ for some $x \in \langle Y_{i-1} \rangle$. Then $[r]_i - 2 \le f_i(z)_i \le [r]_i$.

(b) *For $x, y \in G$, suppose $x \in D_i^r(q)$ and $y \in D_i^s(q)$ for some $1 \le q \le a_i$ and $1 \le r, s \le P_i$. Let $z' = (I_i \circ f_i)(x)$ and $z'' = (I_i \circ f_i)(y)$. Then $||\nu_i(z') - \nu_i(z'')|| \le 3$.*

(c) *For $x, y \in G$, let $I_i(f_i(x)) \in S_i^s(G)$ and $I_i(f_i(y)) \in S_i^t(G)$, $1 \le s \le t \le P_i$.*

(1) If $s = t$, then $|f_{i+1}(x)_{i+1} - f_{i+1}(y)_{i+1}| \le 1$.

(2) If $|s - t| = 1$, then $|f_{i+1}(x)_{i+1} - f_{i+1}(y)_{i+1}| \le 2$.

(d) *Suppose $x \in D_i^s$, $y \in D_i^t$, $1 \le s \le t \le P_i$.*

(1) If $s = t$, then $|f_{i+1}(x)_{i+1} - f_{i+1}(y)_{i+1}| \le 2$.

(2) If $|s - t| = 1$, then $|f_{i+1}(x)_{i+1} - f_{i+1}(y)_{i+1}| \le 3$.

Proof. Consider (a). Since $(I_i \circ f_i)(D_i^{(j)}) \subseteq S_i^{(j)}$ by Lemma 5.1f, it suffices to show that $(I_i \circ f_i)(D_i^j) \cap S_i^{(j-2)} = \emptyset$. Recall that $D_i^{(j-1)} = D_{i-1}^{(a_i(j-1))}$. By Lemma 5.1e1 we have $[a_i(j-1)]_i = \lceil \frac{|D_{i-1}^{(a_i(j-1))}|}{2^{e_{i-1}}} \rceil = \lceil \frac{(j-1)a_i a_{i-1} \cdots a_1}{2^{e_{i-1}}} \rceil$. So by Lemma 4.1 we see that $[a_i(j-1)]_i$ is the number of nonblank $(i-1)$-levels in $S_i^{(j-1)}$. Since I_i preserves the order (by increasing i coordinate) of $(i-1)$-levels, it follows that $I_i(Y_i^{([a_i(j-1)]_i)}) = S_i^{(j-1)}(G)$. By Theorem 6.1b, only the top two $(i-1)$-levels of $Y_i^{([a_i(j-1)]_i)}$ may contain points not in $f_i(D_{i-1}^{(a_i(j-1))}) = f_i(D_i^{(j-1)})$. Again since I_i preserves the order of $(i-1)$-levels, it follows that only the top two nonblank $(i-1)$-levels of $I_i(Y_i^{([a_i(j-1)]_i)}) = S_i^{(j-1)}$, call them Y_i^c and Y_i^d, may contain points not in $(I_i \circ f_i)(D_i^{(j-1)})$. Thus $(I_i \circ f_i)(D_i^j) \cap S_i^{(j-1)} \subset (Y_i^c \cup Y_i^d)$. But since $a_i \ge 2^{22} > 4$ for all i, these top two nonblank

$(i-1)$-levels Y_i^c, Y_i^d of $S_i^{(j-1)}$ lie in the last i-section S_i^{j-1} of $S_i^{(j-1)}$. So $(I_i \circ f_i)(D_i^j) \cap S_i^{(j-2)} = \emptyset$ as desired.

We now consider the consequences (a1)-(a3) of (a), starting with (a1). By (a) we have $(I_{i-1} \circ f_{i-1})(D_{i-1}^r) \subseteq S_{i-1}^{r-1} \cup S_i^r$. Thus $f_i(D_{i-1}^r) \subseteq \sigma_{i-1}(S_{i-1}^{r-1}(G)) \cup \sigma_{i-1}(S_{i-1}^r(G))$. Also the projection map $\bar{\sigma}_{i-1}$ is one to one when restricted to any single $(i-1)$-section, in particular to $S_{i-1}^{r-1}(G)$ and to $S_{i-1}^r(G)$. Thus $f_i(D_{i-1}^r)$ can contribute at most two points to any single stack $Stack_i(x,r)$; namely, up to one point from each of $S_{i-1}^{r-1}(G)$ and $S_{i-1}^r(G)$, proving (a1).

Part (a2) follows immediately from page monotonicity of $Stack_i'(x,r)$ (Theorem 6.1(a)).

Consider (a3). For the upper bound note that $(I_{i-1} \circ f_{i-1})(D_{i-1}^r) \subset S_{i-1}^{(r)}$ by part (a). Since $z \in D_{i-1}^r$, we have $f_i(z)_i \leq \max\{\sigma_{i-1}(u)_i : u \in S_{i-1}^{(r)}(G)\} = [r]_i$ by definition of $[r]_i$. Consider now the lower bound in (a3). If $f_i(z)$ is the highest point in $Stack_i'(x,r)$, then (a3) follows immediately from Theorem 6.1b. Otherwise, by part (a2) we have that $f_i(z)$ is the second highest point in $Stack_i'(x,r)$. Let the highest such point point be $f_i(y)$, $y \in D_{i-1}^r$, so that $f_i(y)_i - f_i(z)_i = 1$ again by part (a2). As in the proof of (a1), we must have $(I_{i-1} \circ f_{i-1})(z) \in S_{i-1}^{r-1}$ and $(I_{i-1} \circ f_{i-1})(y) \in S_{i-1}^r$. Thus by stack monotonicity (Lemma 5.1g) $f_i(z)$ is the highest point in $Stack_i(x, r-1)$, so $f_i(z)_i = |Stack_i(x, r-1)|$. Since $r-1 < P_{i-1}$, by Lemma 5.1d2 we have $f_i(z)_i = |Stack_i(x, r-1)| \geq [r-1]_i - 1$. It remains to check that $[r-1]_i - 1 \geq [r]_i - 2$, which follows directly from the formula $[r]_i = \lceil \frac{r a_{i-1} a_{i-2} \cdots a_1}{2^{e_{i-1}}} \rceil$ proved in Lemma 5.1e1.

For (b), recall that $D_i^r(q) = D_{i-1}^{(r-1)a_i+q}$, with the same formula for $D_i^s(q)$ where r is replaced by s. Let $d_i = a_i a_{i-1} \ldots a_1$. By part (a) we have $(I_{i-1} \circ f_{i-1})(D_{i-1}^{(r-1)a_i+q}) \subseteq S_{i-1}^{(r-1)a_i+q-1} \cup S_{i-1}^{(r-1)a_i+q}$. Hence using Lemma 5.1e1 and the upper bound in part (a3), we have $f_i(x)_i \leq [(r-1)a_i+q]_i = \lceil \frac{((r-1)a_i+q)d_{i-1}}{2^{e_{i-1}}} \rceil$. Now by Lemma 4.1a and our construction we know that $S_i^{(r-1)}$ has $\lceil \frac{(r-1)d_i}{2^{e_{i-1}}} \rceil = \lceil \frac{(r-1)a_i d_{i-1}}{2^{e_{i-1}}} \rceil$ nonblank $(i-1)$-levels. Thus the first $\lceil \frac{(r-1)a_i d_{i-1}}{2^{e_{i-1}}} \rceil$ nonblank blank $(i-1)$-levels in the image $I_i(Y_i^{([(r-1)a_i+q]_i)})$ belong to $S_i^{(r-1)}$. Hence $\nu_i(z') \leq \lceil \frac{((r-1)a_i+q)d_{i-1}}{2^{e_{i-1}}} \rceil - \lceil \frac{(r-1)a_i d_{i-1}}{2^{e_{i-1}}} \rceil$. Similarly, this time using Lemma 5.1e1 and the lower bound in (a3), we have $\nu_i(z'') \geq \lceil \frac{((s-1)a_i+q)d_{i-1}}{2^{e_{i-1}}} \rceil - 2 - \lceil \frac{(s-1)a_i d_{i-1}}{2^{e_{i-1}}} \rceil$. It follows that $\nu_i(z') - \nu_i(z'') \leq 3$. A symmetric argument interchanging the roles of r and s yields $\nu_i(z') - \nu_i(z'') \geq -3$, completing (c).

Consider part (c), starting with (c1). Let $x' \in \langle Y_i \rangle$ and $y' \in \langle Y_i \rangle$ be the stack addresses of the images $f_{i+1}(x)$ and $f_{i+1}(y)$; that is $x' = f_{i+1}(x)_{1 \to i}$ and $y' = f_{i+1}(x)_{1 \to i}$. Since $s-1 \leq P_i - 1$, by Lemma 5.1d we have $[s-1]_{i+1} - 1 \leq |Stack_{i+1}(x', s-1)|, |Stack_{+1}(y', s-1)| \leq [s-1]_{i+1}$. Since $I_i(f_i(x)) \in S_i^s(G)$, we have $f_{i+1}(x) = \sigma_i(I_i(f_i(x))) \in Stack_{i+1}(x', s) - Stack_{i+1}(x', s-1)$. By Lemma 5.1c $f_{i+1}(x)$ is the only member of $Stack_{i+1}(x', s) - Stack_{i+1}(x', s-1)$, and thus $|Stack_{i+1}(x', s)| \leq |Stack_{i+1}(x', s-1)| + 1$. It follows that $[s-1]_{i+1} \leq f_{i+1}(x)_{i+1} \leq [s-1]_{i+1} + 1$, with the same inequality holding for $f_{i+1}(y)_{i+1}$. Part (c1) follows.

For (c2), suppose without loss of generality that $s = t-1$. Again using Lemma 5.1c, we get $[s]_{i+1} \leq [t]_{i+1} \leq [s]_{i+1} + 1$. Since $I_i(f_i(y)) \in S_i^t(G)$ we have $f_{i+1}(y)_{i+1} \leq [t]_{i+1}$ by definition of $[t]_{i+1}$. Again since $I_i(f_i(y)) \in S_i^t(G) = S_i^{s+1}(G)$ and using Lemma 5.1d2 we have $f_{i+1}(y)_{i+1} = 1 + |Stack_{i+1}(y', s)| \geq 1 + [s]_{i+1} - 1 = [s]_{i+1}$. Hence we have $[s]_{i+1} \leq f_{i+1}(y)_{i+1} \leq [s]_{i+1} + 1$. Since $s \leq P_i - 1$ and $f_{i+1}(x)$ is the topmost entry of $Stack_{i+1}(x', s)$, it follows by Lemma 5.1d2 that $[s]_{i+1} - 1 \leq f_{i+1}(x)_{i+1} \leq [s]_{i+1}$. Part (c2) follows.

For (d), set $\alpha = I_i(f_i(x))$ and $\beta = I_i(f_i(y))$, and consider first (d1). By part (a), α and β lie in the same i-section or in successive i-sections S_i^{s-1}, S_i^s of Y_i. Hence (d1) follows directly from part (c). For (d2), suppose $s = t-1$. Again by part (a), $\alpha \in S_i^{s-1} \cup S_i^s = S_i^{t-2} \cup S_i^{t-1}$,

and $\beta \in S_i^{t-1} \cup S_i^t$. If α and β belong to successive i-sections (e.g. S_i^{t-2} and S_i^{t-1}, or S_i^{t-1} and S_i^t respectively), then $|f_{i+1}(x)_{i+1} - f_{i+1}(y)_{i+1}| \leq 2$ by part (c). So we can suppose that $\alpha \in S_i^{t-2}$ and $\beta \in S_i^t$. Since the projection $\bar{\sigma}_i$ is one to one on any one i-section (Lemma 5.1c), we have $[t]_{i+1} \leq [t-2]_{i+1} + 2$. Since $t - 1 \leq P_i - 1$ we have $f_{i+1}(x)_{i+1} \geq [t-2]_{i+1} - 1$ by Lemma 5.1d, while $f_{i+1}(y)_{i+1} \leq [t]_{i+1}$ since $\beta \in S_i^t$. Thus $|f_{i+1}(x)_{i+1} - f_{i+1}(y)_{i+1}| \leq 3$, completing (d2). ■

We now proceed to a bound on the dilation of our embedding. In the proof we frequently apply Corollary 4.5, where (in the language of its statement) we use $N_u(d)$ relative to the matrix $F^X = (f_{uv})$ with the settings in part (d) of that Corollary. By the construction given immediately after that Corollary, we have $F^X = F(i)$. The value of i will change from one application to another. So in the proof which follows, let $N_{i,u}(d)$ denote the $N_u(d)$ of the corollary, when in the application we intend to use $F^X = F(i)$. So for example, in applying Corollary 4.5 with $e = 3$ and $F^X = F(2)$, we would conclude that $|N_{2,u}(d+3) - N_{2,u}(d)| \leq 2 \cdot 3 + 2 = 8$, using the more generous of the bounds (a) and (b).

Theorem 6.3 *Let $G = [a_1 \times a_2 \times \cdots \times a_k]$ be a k-dimensional grid with $a_i \geq 2^{22}$ for each i. Then the embedding $H^k : G \to Opt(G)$ from section 5.2 satisfies dilation$(H^k) \leq 3k$.*

Proof. Let $x = (x_1, x_2, \ldots, x_k)$ and $y = (y_1, y_2, \ldots, y_k)$ be arbitrary adjacent points of G, and set $Q = Opt(G)$ and $f = f_k$ for short. Recall that
$H^k(x) = (H^k(x)_1, H^k(x)_2, \ldots, H^k(x)_j, \ldots, H^k(x)_k) \in Q$, where $H^k(x)_j = L^{-1}_{e_j - e_{j-1}}(f(x)_j)$ is the $(0,1)$ string of $Q_{e_j - e_{j-1}}$ equivalent to $f(x)_j$ under the labeling $L_{e_j - e_{j-1}}$ of Corollary 4.8. It suffices to show for each j, $1 \leq j \leq k$, that $|f(x)_j - f(y)_j| \leq 17$. For then by the definition of H^k just given and Corollary 4.8b we have $dist_{Q_{e_j - e_{j-1}}}(H^k(x)_j, H^k(y)_j) \leq 3$ for each $1 \leq j \leq k$. So we obtain $dist_Q(H^k(x), H^k(y)) = \sum_{j=1}^k dist_{Q_{e_j - e_{j-1}}}(H^k(x)_j, H^k(y)_j) \leq 3k$, and our desired dilation bound follows. Thus we are reduced to showing that $|f(x)_j - f(y)_j| \leq 17$ for each j.

Let $i_0 := i_0(x,y)$, $1 \leq i_0 \leq k$, be the unique coordinate at which the above adjacent points $x, y \in V(G)$ disagree, so $|x_{i_0} - y_{i_0}| = 1$. Also let $L(j, i_0) = max\{|f(x)_j - f(y)_j| : xy \in E(G), |x_{i_0} - y_{i_0}| = 1\}$. For any given j, the maximum of $L(j, i_0)$ over all i_0 serves as an upper bound for $|f(x)_j - f(y)_j|$, over all edges $xy \in E(G)$. So it suffices to show that that for each $1 \leq j \leq k$ this maximum is at most 17. Let $x(j)y(j)$ be an edge at which this maximum occurs for a given j, and refer to this edge simply as xy since j will be understood by context.

We begin with $j = 1$, starting with the case $i_0 = 1$ for a bound on $L(1,1)$. Since $i_0 = 1$, x and y are corresponding points on successive chains C_i and C_{i+1} of $G(a_1)$. So we have $L(1,1) = |f(x)_1 - f(y)_1| = |f_2(x)_1 - f_2(y)_1| \leq 3 < 17$ as required, where the second equality follows from Lemma 5.1a and the inequality following that from Corollary 2.3a. Now suppose $i_0 \geq 2$. Thus x and y agree in their first coordinate, so can be considered as lying on the same chain of $G(a_1)$. So similarly by by Lemma 5.1a and Theorem 2.2h we have $L(1, i_0) = |f(x)_1 - f(y)_1| = |f_2(x)_1 - f_2(y)_1| \leq 2 < 17$. So $\max\{L(1, i_0) : 1 \leq i_0 \leq k\} \leq 17$.

Now suppose that $j = 2$. If $i_0 = 1$, then again x and y are corresponding points on successive chains of $G(a_1)$. Thus $|f_2(x)_2 - f_2(y)_2| \leq 1$ by Theorem 2.2g. Hence by Lemma 5.1b (with $e = 1$) we get $L(2,1) = |f(x)_2 - f(y)_2| \leq 4 < 17$.

If $i_0 = 2$, then x and y are successive points on the same chain of $G(a_1)$, so by Corollary 2.3a we have $|f_2(x)_2 - f_2(y)_2| \leq 1$. So again by Lemma 5.1b, $L(2,2) = |f(x)_2 - f(y)_2| \leq 4 < 17$.

Now suppose $i_0 \geq 3$, still with $j = 2$. Then x and y belong to the same chain C_i of $G(a_1)$. Since $x_1 = y_1$, $x_2 = y_2$, and $i_0 \geq 3$, we can view x and y as corresponding points belonging to a pair of distinct 2-pages of G, say $x \in D_2^u$ and $y \in D_2^v$. Consider the segments T_1 and T_2 of C_i given by $T_1 = \{(i, (u-1)a_2 + t) : 1 \leq t \leq x_2\}$ and $T_2 = \{(i, (v-1)a_2 + t) : 1 \leq t \leq x_2\}$. We see that $f_2(T_1)$ (resp. $f_2(T_2)$) is the initial segment of x_2 points of the chain C_i in D_2^u (resp. D_2^v), and that $f_2(x)$ and $f_2(y)$ are the last points of these segments respectively. Let c_1 and c_2 be the number of successive columns of Y_2 spanned by $f_2(T_1)$ and $f_2(T_2)$ respectively. By Corollary 2.3d we have $|c_1 - c_2| \leq 1$. The columns spanned by $f_2(T_1)$ (resp. $f_2(T_2)$) are transformed under the inflation map I_2 into a corresponding set of c_1 (resp. c_2) successive nonblank columns in $(I_2 \circ f_2)(G)$. Corollary 2.3e and Lemma 4.1 now imply that $(I_2 \circ f_2)(T_1)$ and $(I_2 \circ f_2)(T_2)$, being transforms of $f_2(T_1)$ and $f_2(T_2)$ respectively, each begin in either the first nonblank column of their section (S_2^u for $(I_2 \circ f_2)(T_1)$ and S_2^v for $(I_2 \circ f_2)(T_2)$) or the last nonblank column of their preceding section (S_2^{u-1} for $(I_2 \circ f_2)(T_1)$ and S_2^{v-1} for $(I_2 \circ f_2)(T_2)$). Now let $z' = (I_2 \circ f_2)(x)$ and $z'' = (I_2 \circ f_2)(y)$, and assume by symmetry that $c_2 \geq c_1$. Then $|c_1 - c_2| \leq 1$ implies that $\nu_2(z'') - \nu_2(z') = 0, 1$ or 2. Assume first that $\nu_2(z'') - \nu_2(z') = 2$, and write $d = \nu_2(z')$ and $d+2 = \nu_2(z'')$. This case arises when $c_2 - c_1 = 1$, and $(I_2 \circ f_2)(T_2)$ begins in the first nonblank column of S_2^v while $(I_2 \circ f_2)(T_1)$ begins in the last nonblank column of S_2^{u-1}. Then by Lemma 5.1a and Corollary 4.5c with $e = 2$, we get $|f(x)_2 - f(y)_2| = |f_3(x)_2 - f_3(y)_2| = |N_{2,u}(\nu_2(z')) - N_{2,v}(\nu_2(z''))| = |N_{2,u}(d) - N_{2,v}(d+2)| \leq 2 \cdot 2 + 4 = 8$. In the case $\nu_2(z'') - \nu_2(z') = 1$, we similarly obtain (using $e = 1$ in Corollary 4.5c) that $|f(x)_2 - f(y)_2| \leq 6$, and in the case $\nu_2(z'') - \nu_2(z') = 0$ (using $e = 0$) we get $|f(x)_2 - f(y)_2| \leq 4$. So we get $L(2, i_0) \leq 8 < 17$ for $i_0 \geq 3$, and overall $\max\{L(2, i_0) : 1 \leq i_0 \leq k\} \leq 17$. .

Next suppose $j \geq 3$. We argue according to the order relation between j and i_0.

Suppose first that $j > i_0$. Then x and y belong to the same $(j-1)$-page and successive $(i_0 - 1)$-pages of G. We obtain our bound on $L(j, i_0)$ independent of i_0. Since x and y belong to the same $(j-1)$-page we have that $I_{j-1}(f_{j-1}(x))$ and $I_{j-1}(f_{j-1}(y))$ belong to the same $(j-1)$-section or to successive $(j-1)$-sections S_{j-1}^t and $S_{j-1}^{t'}$, $|t - t'| \leq 1$, by Corollary 6.2a. Applying Corollary 6.2c, in the first case (same $(j-1)$-section) we get $|f_j(x)_j - f_j(y)_j| \leq 1$ by part (c1), while in the second case (successive $(j-1)$-sections) we get $|f_j(x)_j - f_j(y)_j| \leq 2$ by part (c2). Now we bound $L(j, i_0)$ by applying Lemma 5.1b (with $e = 2$) to obtain $L(j, i_0) = |f(x)_j - f(y)_j| \leq 6 < 17$, as required.

Now suppose $j = i_0$. Then x and y are corresponding points in successive $(j-1)$-pages of G. Hence by Corollary 6.2d2 we get $|f_j(x)_j - f_j(y)_j| \leq 3$. Thus by Lemma 5.1b with $e = 3$ we get $L(j, j) = |f(x)_j - f(y)_j| \leq 8 < 17$.

It remains to consider the case $3 \leq j < i_0$. Here x and y are corresponding points in successive $(i_0 - 1)$-pages, say $x \in D_{i_0-1}^r$ and $y \in D_{i_0-1}^{r+1}$. Consider the j-pages containing x and y, say $x \in D_j^c \subset D_{i_0-1}^r$ and $y \in D_j^d \subset D_{i_0-1}^{r+1}$ for suitable integers c and d. Since x and y agree in their first j coordinates, they must also be corresponding points in these j-pages. In particular, x and y belong to corresponding $(j-1)$-subpages $D_j^c(q)$ of D_j^c and $D_j^d(q)$ of D_j^d respectively for the same integer q. As before, let $z' = (I_j \circ f_j)(x)$ and $z'' = (I_j \circ f_j)(y)$ for brevity. Then by Corollary 6.2b we get $|\nu_j(z') - \nu_j(z'')| \leq 3$. Again applying Lemma 5.1a we have $f(x)_j = N_{j,c}(\nu_j(z'))$ and $f(y)_j = N_{j,d}(\nu_j(z''))$. So applying Corollary 4.5c with $e = 3$ we get $L(j, i_0) = |f(x)_j - f(y)_j| \leq |N_{j,c}(\nu_j(z')) - N_{j,d}(\nu_j(z''))| \leq 2 \cdot 3 + 4 = 10 < 17$. So for any $j \geq 3$ we have $\max\{L(j, i_0) : 1 \leq i_0 \leq k\} \leq 17$, completing the proof. ∎

7 Concluding Remarks

1. There is a routing of edge congestion $O(k)$ associated to our embedding H^k. We outline the idea here, omitting full details of the proof.

As notation, for any graph H and permutation $\pi : V(H) \to V(H)$, a *π-routing* is an assignment $P : V(H) \to \{\text{paths in } H\}$ such that $P(x)$ is a path in H from x to $\pi(x)$. If H is directed, then $P(x)$ is a directed path from x to $\pi(x)$. The *congestion* of a π-routing is the maximum of $\{n(e) : e \in E(H)\}$, where $n(e)$ is the number of paths in the π-routing which use the edge e.

For background, let $\overrightarrow{Q_n}$ be the directed graph obtained from Q_n by replacing each edge of Q_n by 4 directed edges, two pointing in one direction and two in the opposite direction. Using the classic Benes routing method, one can show [14] that for any permutation π of $\overrightarrow{Q_n}$ there exists a π-routing such that the paths of the routing are edge disjoint. Consequently, for each permutation π of the undirected Q_n there is a π-routing with congestion $O(1)$.

For an undirected graph H, suppose there is a partition of $E(H)$ into k sets, $E(H) = \bigcup_{i=1}^{k} E_i$, such that each E_i is a vertex disjoint union of cycles in H, where the vertices of each cycle are ordered in one of the two natural ways. Further, let $g : H \to Q_n$ be a one to one map. Now consider the permutation π_i on Q_n as follows: if $v \notin g(H)$ then $\pi_i(v) = v$, while if $v \in g(H)$ then $\pi_i(v) = w$, where $g^{-1}(v)g^{-1}(w) \in E_i$ and $g^{-1}(w)$ follows $g^{-1}(v)$ in the ordering of the vertices of the cycle of E_i containing $g^{-1}(v)$ and $g^{-1}(w)$. Then by the above result there is a π_i routing on Q_n of congestion $O(1)$ for each i, $1 \le i \le k$. Since each $e \in E(H)$ lies in some E_i, it would follow that the edge congestion of g (as defined in section 1.1) is $O(k)$.

It therefore suffices to find a graph $G' \supset G = [a_1 \times a_2 \times \ldots \times a_k]$ with $V(G') = V(G)$ such that G' has the required cycle partition of edges. For then (letting our map H^k play the role of g in the above paragraph) the map $H^k : G' \to Opt(G)$ has edge congestion $O(k)$ by the above argument, so the same is true of its restriction $H^k : G \to Opt(G)$.

The graph G' is obtained from G as follows. For each i, $1 \le i \le k$, consider a fixed $(k-1)$-tuple $\mathbf{c}(i) = (c_1, c_2, \cdots, c_{i-1}, c_{i+1}, \cdots, c_k)$ of integer entries, with $1 \le c_j \le a_j$ for $j \ne i$. Further, for $1 \le t \le a_i$ let $v(\mathbf{c}(i), t) = (c_1, c_2, \cdots, c_{i-1}, t, c_{i+1}, \cdots, c_k) \in V(G)$. We let $V(G') = V(G)$ and $E(G') = E(G) \cup E'$, where $E' = \{v(\mathbf{c}(i), a_i)v(\mathbf{c}(i), 1) : 1 \le i \le k, \mathbf{c}(i) \text{ any } (k-1)\text{-tuple as above}\}$. Thus E' is just the set of "wraparound" edges not present in G in each of the k dimensions. The required cycle partition of $E(G')$ is given by $E(G') = \bigcup_{i=1}^{k} E'_i$, where $E'_i = \{v(\mathbf{c}(i), a_i)v(\mathbf{c}(i), 1), v(\mathbf{c}(i), t)v(\mathbf{c}(i), t+1) : 1 \le t \le a_i - 1, 1 \le i \le k, \mathbf{c}(i) \text{ any } (k-1)\text{-tuple as above}\}$.

2. The lower bound requirement $a_i > 2^{22}$ for our result can be relaxed to $a_i > 2^{12}$, provided one can improve the conclusion of Corollary 4.8a only slightly to say $|L_t(x) - L_t(y)| \le 2r+4 \Rightarrow dist_{Q_t}(x, y) \le 3$. Then using $r = 3$ and $i = 2$ we obtain $|L_t(x) - L_t(y)| \le 10 \Rightarrow dist_{Q_t}(x, y) \le 3$ for $t \ge 12$, so that $a_i > 2^{12}$ suffices for our result. The proof of Theorem 6.3 is then reduced (as in its first paragraph) to proving the inequality $|f(x)_j - f(y)_j| \le 10$ for each $1 \le j \le k$ and $xy \in E(G)$. The proof of Theorem 6.3 shows that this inequality does indeed hold. The improvement in Corollary 4.8a may require a detailed study of the regular cyclic caterpillars we used.

3. The question of finding good lower bounds for $B(G, Opt(G))$, for some class of multidimensional grids G with $|V(G)| \to \infty$, remains open. A nontrivial lower bound for all multidimensional grids is of course not possible, since if the a_i are all powers of 2, then G is

a spanning subgraph of $Opt(G)$, so $B(G, Opt(G)) = 1$ in that case.

4. The first author thanks Stephen Wright for many useful discussions on rounding of matrices. We also acknowledge a useful discussion with Hal Kierstead which inspired our application of regular cyclic caterpillars.

References

[1] T. Asano, T. Matsui, and T. Tokuyama: Optimal Roundings of Sequences and Matrices, *Nordic J. Comput.* **7** (2000) 241-256.

[2] T. Asano, N. Katoh, K. Obokata, and T. Tokuyama: Matrix Rounding Under the L_p-Discrepancy Measure and Its Application to Digital Halftoning, *SIAM J. Comput.* **32** No.6 (2003) 1423-1435.

[3] U. Blass: Short Dominating Paths and Cycles in the Binary Hypercube: *Annals of Combinatorics* **5** No. 1 (2001) 51-59.

[4] Zs. Baranyai: On the Factorization of the Complete Uniform Hypergraph, in *Infinite and Finite Sets, Proc. Coll. Keszthely, 1975, Colloquia Math. Soc. János Bolyai* A. Hajnal, R. Rado, V.T. Sós (Eds.), Colloquia Math. Soc. János Bolyai **10** North-Holland (1975) 91-108.

[5] S. Bettayeb, Z. Miller, and I.H. Sudborough: Embedding Grids into Hypercubes, *J. Comput. System Sci.* **45** No. 3 (1992) 340-366.

[6] S. Bhatt, F.R.K. Chung, F.T. Leighton, and A. Rosenberg: Efficient Embeddings of Trees in Hypercubes, *SIAM J. Comput.* **21** No.1 (1992) 151-162.

[7] R. Caha and V. Koubek: Spanning Regular Caterpillars in Hypercubes, *European J. Combinatorics* **18** (1997) 249-266.

[8] Y. Cai, T.H. Lai, and C.C. Lin: Embedding Two-Dimensional Grids into Hypercubes with Dilation 2 and Congestion 5, Proc. Intern. Conf. on Parallel Processing, Oconomowoc, Wisc., III (1995) 105-112.

[9] M.Y. Chan: Embeddings of Grids into Optimal Hypercubes, *SIAM J. Comput.* **20** No. 5 (1991) 834-864.

[10] B. Doerr: Matrix Rounding with Respect to Small Submatrices, *Random Structures & Algorithms* **28** (2006) 107-112.

[11] B. Doerr, T. Friedrich, C. Klein, R. Osbild: Rounding of Sequences and Matrices, with Applications, *Proceedings of the Workshop on Approximation and Online Algorithms (WAOA 2005)* T. Erlebach, G. Persiano (Eds.) Vol. 3879 of Lecture Notes in Computer Science, Springer Verlag (2006) 96-109.

[12] B. Doerr, T. Friedrich, C. Klein, and R. Osbild: Unbiased Matrix Rounding, *Proceedings of the 10th Scandinavian Workshop on Algorithm Theory (SWAT)*, L. Arge, R. Freivalds (Eds.) Vol. 4059 of Lecture Notes in Computer Science, Springer Verlag (2006) 102-112.

[13] T. Dvorak, I. Havel, J-M Laborde, and M. Mollard: Spanning Caterpillars of a Hypercube, *Journal of Graph Theory* **24** No. 1 (1997) 9-19.

[14] Q.-P. Gu and H. Tamaki: Routing a Permutation in the Hypercube by Two Sets of Edge Disjoint Paths, *J. Parallel Distrib. Comput.* **44**(2) (1997) 147-152.

[15] I. Havel and P. Liebl: Embedding the Polytomic Tree into the n-cube, *Casopis Pest. Mat.* **98** (1973) 307-314.

[16] I. Havel and J. Moravek: B-Valuations of Graphs, *Czech. Math. J.* **22** (1972) 338-351.

[17] F. Harary, M. Lewinter, and W. Widulski: On Two-Legged Caterpillars which Span Hypercubes *Congress. Numer.* (19'th Southeastern Conference on Comb., Graph Theory, and Computing) **66** (1988) 103-108.

[18] Y.M. Kim and T.H. Lai: The Complexity of Congestion-1 Embedding in a Hypercube, *J. of Algorithms* **12** No. 2 (1991) 246-280.

[19] D.E. Knuth: Two-Way Rounding, *SIAM J. Disc. Math* **8** No. 2 (1995) 281-290.

[20] J. Lee: On the Efficient Simulation of Networks by Hypercube Machines, Ph.D. Dissertation, Dept. of Electrical Engineering and Computer Science, Northwestern University (1990).

[21] F.T. Leighton: *Introduction to Parallel Algorithms and Architectures: Arrays, Trees, Hypercubes*, Morgan Kaufman Publishers, San Mateo, CA (1992).

[22] M. Livingston and Q. Stout: Embeddings in Hypercubes, *Mathematics and Computational Modelling* **11** (1988) 222-227.

[23] L. Lovasz, J. Spencer, and K. Vestergombi: Discrepancy of Set Systems and Matrices, *European J. of Combin.* **7** (1986) 151-160.

[24] Z. Miller and I.H. Sudborough: Compressing Grids into small Hypercubes, *Networks* **24** No. 6 (1994) 327-357.

[25] A. L. Rosenberg, L. S. Heath: *Graph Separators with Applications*, Kluwer Academic/Plenum Publishers, New York, New York (2001).

[26] M. Rottger, U. Schroeder: Embedding 2-Dimensional Grids into Optimal Hypercubes with Edge Congestion 1 or 2, *Parallel Processing Letters* **8** No. 2 (1998) 231-242.

[27] J. Spencer: Ten Lectures on the Probabilistic Method: *CBMS-NSF Regional Conference Series in Applied Mathematics* **52** SIAM, Philadelphia (1987) 37-44.

[28] D.B. West: *Introduction to Graph Theory* (Second Edition) Prentice Hall (2001).

[29] S. E. Wright: Integer Matrices with Constraints on Leading Partial Row and Column Sums, *Discrete Applied Mathematics* Vol. 158 Issue 16 (2010) 1838-1847.

8 Appendix 1: Proofs of Theorem 2.2 and Corollary 2.3

Proof of Theorem 2.2: Part (a) follows directly from steps 2a and 2b of the construction of f. In part (b), the claim that $L_r(j)$ is an initial segment of Y_2^j follows directly from part (a), step 2 of the construction of f, and induction on r. The claim on $|L_r(j)|$ follows from step 2 of this construction on noting that C_i contributes 1 (resp. 2) to this sum if $R_{ij} = 0$ (resp. $R_{ij} = 1$).

Consider (c). By part (a), $\sum_{t=1}^{j} R_{it}$ is the number of columns Y_2^t, $1 \le t \le j$, such that $|f(C_i) \cap Y_2^t| = 2$, while in the remaining columns Y_2^t, $1 \le t \le j$, we have $|f(C_i) \cap Y_2^t| = 1$. So $\pi(i, 1, j) = N_{ij}$ as claimed. Also observe that $\pi(i, r \to r+j) = N_{i,r+j} - N_{i,r-1} = j + 1 + \sum_{t=r}^{r+j} R_{it} \in \{j + 1 + S_{j+1}, j + 2 + S_{j+1}\}$ by the formula for N_{ij} and Theorem 2.1b. The same equation with r replaced by s yields the rest of (c).

Part (d) follows from parts (b), (c), Lemma 2.1b, and the fact that $S_{r+1} - S_r \le 1$. For (e), by part (c) (taking $j = m$) and the definition of $G(a_1, N_{im})$ we see that $f(G(a_1, N_{im})) \subseteq Y_2^{(m)}$. So it suffices to show that $|f(G(a_1, N_{im}))| = |Y_2^{(m)}|$. We have $|f(G(a_1, N_{im}))| = \sum_{i=1}^{a_1} \sum_{j=1}^{m} N_{ij} = \sum_{i=1}^{a_1} (m + \sum_{t=1}^{m} R_{it}) = a_1 m + m(2^{L_1} - a_1) = m2^{L_1} = |Y_2^{(m)}|$, using Lemma 2.1a.

Consider (f). Recall the notation $P_1 = a_2 a_3 \cdots a_k$ for the number of 1-pages in G. We use the fact that G can be identified with the the subgraph of $G(a_1)$ induced by the union of initial segments $\bigcup_{i=1}^{a_1} C_i(P_1)$ via the map κ of section 2. For $j = m-1$ or m let $M'_j = min\{N_{ij} : 1 \le i \le a_1\}$, and $M''_j = max\{N_{ij} : 1 \le i \le a_1\}$. We claim that $M'_m \ge P_1$. If not, then $M'_m < P_1$, so $M''_m \le P_1$ by (d). It follows that $|Y_2^{(m)}| = |f(G(a_1, N_{im}))| = \sum_{i=1}^{a_1} N_{im} < a_1 P_1 = |G|$. But this contradicts $|Y_2^{(m)}| = m2^{e_1} \ge |G|$, proving that $M'_m \ge P_1$. Hence for each $1 \le i \le a_1$ we have $N_{im} \ge P_1$, implying that $G \subseteq G(a_1, N_{im})$ by the identification of G above. To show $Y_2^{(m-1)} \subset f(G)$, observe from the definition of m that $|Y_2^{(m-1)}| < |G|$. Thus $\sum_{i=1}^{a_1} N_{i,m-1} = |Y_2^{(m-1)}| < |G| = a_1 P_1$. So one of the terms in the last sum is less than P_1. It follows that $M'_{m-1} < P_1$ and hence $M''_{m-1} \le P_1$ by (d). This says that for each i, $f(i, P_1 + 1)_2 \ge m$, and hence that $f(i, j)_2 \ge m$ for $j \ge P_1 + 1$ by the monotonicity of f from part (a). It follows that $Y_2^{(m-1)} \subset f(\bigcup_{i=1}^{a_1} C_i(P_1)) = f(G)$, as required.

For (g), we induct on r. The base case is clear since $f(i, 1)_2 = 1$ for all i. Assume inductively that $|f(i, r)_2 - f(j, r)_2| \le 1$ for some $r > 1$, and let $c = f(i, r)_2$. If $f(j, r)_2 = f(i, r)_2 = c$, then we are done since $f(i, r+1)_2 = c$ or $c+1$ and the same holds for $f(j, r+1)_2$. So suppose $|f(i, r)_2 - f(j, r)_2| = 1$, and without loss of generality that $f(j, r)_2 = c+1$ (the case $f(j, r)_2 = c-1$ being symmetric). Then $N_{j,c} \le r - 1$ by (c) and $\sum_{t=1}^{c} R_{it} = 1 + \sum_{t=1}^{c} R_{jt}$ by Lemma 2.1. If $f(j, r+1)_2 = f(j, r)_2 = c+1$, then we are done since $f(i, r)_2 \le f(i, r+1)_2 \le f(i, r)_2 + 1$. So we may assume that $f(j, r+1)_2 = c + 2$. If now $f(i, r+1)_2 = c + 1$, then we are done again. So we can also assume that $f(i, r+1)_2 = c$. Then we get $N_{ic} \ge r + 1 \ge N_{jc} + 2$, contradicting $\sum_{t=1}^{c} R_{it} = 1 + \sum_{t=1}^{c} R_{jt}$.

Consider (h). Set $c = f(i, y)_2$ for arbitrary y, $1 \le y \le N_{im}$. It suffices to show that $f(i, y)_1$ must be one of three consecutive integers which depend only on i, but not on y. By (b), the set $L_{i-1}(c)$ is the initial segment of size $i - 1 + \sum_{t=1}^{i-1} R_{tc}$ in Y_2^c. So by Lemma 2.1, $|L_{i-1}(c)|$ is

either $i-1+S_{i-1}$ or $i+S_{i-1}$. Now $f(C_i) \cap Y_2^c$ consists of either one or two successive points of Y_2^c which immediately follow the point $(|L_{i-1}(c)|, c)$. Hence $f(i,y)$ must be one of the three successive integers $i+S_{i-1}, i+1+S_{i-1}$, or $i+2+S_{i-1}$, proving (h).

Finally consider (i). The proof is based on R^{j+1} being a downward shift of R^j (with wraparound) for any j. We refer to this property as the "downward shift" property. The assumption $|f(C_r) \cap Y_2^j| = 2$ says that $R_{rj} = R_{r-j+1,1} = 1$ (viewing subscripts modulo a_1), by the downward shift property. Hence by part (b) and this same property, we have $|L_r(j)| - |L_r(j+1)| = R_{rj} - R_{1,j+1} \geq 0$, since $R_{rj} = 1$. For the second claim, observe that $N_{r,j} = j + \sum_{i=1}^{j} R_{ri} = j + \sum_{i=r-j+1}^{r} R_{i1}$ by the downward shift property. Similarly we have $N_{r+1,j} = j + \sum_{i=r-j+2}^{r+1} R_{i1}$. Therefore $N_{r,j} - N_{r+1,j} = R_{r-j+1,1} - R_{r+1,1} \geq 0$, since $R_{r-j+1,1} = 1$.

This completes the proof of the theorem.

Proof of Corollary 2.3: As a convenience, we prove these properties with f replacing f_2, to facilitate direct reference to the construction above. Of course f_2 then inherits these properties since it is a restriction of f. For brevity let $d_1 = |f(v)_1 - f(w)_1|$ and $d_2 = |f(v)_2 - f(w)_2|$.

Consider part (a). We get $d_2 \leq 1$ by Theorem 2.2a if v and w are successive points on the same chain, and by Theorem 2.2g otherwise (i.e., if $v = (i,j)$ and $w = (i+1,j)$ for some i and j). We also get $d_1 \leq 2$ by Theorem 2.2h if v and w are successive points on the same chain. So we are reduced to showing that $d_1 \leq 3$ when, say, $v = (i,j)$ and $w = (i+1,j)$ for some i and j. Let $f(v)_2 = c$. If also $f(w)_2 = c$ then since by Theorem 2.2a,b we have that $(f(C_i) \cup f(C_{i+1})) \cap Y_2^c$ is a set of at most 4 successive points of Y_2^c, it follows in this case that $d_1 \leq 3$.

So let $f(w)_2 = c' \neq c$, noting that then $|c-c'| = 1$ since we have already shown that $d_2 \leq 1$. By Theorem 2.2a,b we have $f(v)_1 = |L_i(c)|$ or $|L_i(c)|-1$, and similarly $f(w)_1 = |L_{i+1}(c'+1)|$ or $|L_i(c')|-1$. By Theorem 2.2d we have $||L_{i+1}(c')| - |L_i(c)|| \leq 2$, so $d_1 \leq 3$ follows, completing part (a).

The proof of (b) is a tedious case analysis showing that the bound on d_1 from part (a) can be strengthened to $d_1 \leq 2$, which combined with (a) yields (b). The proof is omitted here for brevity. We include (b) only for its possible interest, and do not use it later.

Consider part (c). Since $m = \lceil \frac{|G|}{2^{e_1}} \rceil$, we see that $Y_2^{(m)}$ is a subgraph of the spanning subgraph $Y_2^{(2^{e_k - e_1})}$ of $Opt(G)$. Thus by Theorem 2.2e,f we obtain $f_2(G) \subset f(H) = Y_2^{(m)} \subset Opt(G)$, as required.

Consider next part (d). As notation, for $a \leq b$ let $Y_2^{(a \to b)} = \bigcup_{t=a}^{b} Y_2^t$ be the union of columns a through b of Y_2. Let then $Y_2^{(r \to r+c-1)}$ and $Y_2^{(s \to s+c'-1)}$ be the sets of columns of Y_2 spanned by $f_2(T)$ and $f_2(T')$ respectively. Suppose to the contrary that $|c-c'| \geq 2$. By symmetry we can assume that $c' \geq c+2$.

Let $S = f_2(T) \cap Y_2^{(r \to r+c-1)}$, and let $S' = f_2(T') \cap Y_2^{(s+1 \to s+c)}$. Now for each $t, 1 \leq t \leq s+c'-1$, we have $f_2(T') \cap Y_2^t = C_j \cap Y_2^t$ with the possible exception when $t = s$ (resp. $t = s+c'-1$), where possibly $f_2(T')$ contains just the second (resp. first) of two points of $C_j \cap Y_2^t$. So since $c < c'-1$ it follows that $S' = f_2(C_j) \cap Y_2^{(s+1 \to s+c)}$. So $|S'| = \pi(j, s+1 \to s+c)$. Thus by Theorem 2.2c we have $||S'| - \pi(i, r \to r+c-1)| \leq 1$. Now $|S'| \leq p-2$, since S' omits at least the two endpoints of the path $f_2(T')$. But also $\pi(i, r \to r+c-1) \geq |S| \geq p$, a contradiction.

Finally we prove part (e). The containment in (e) follows from (d) (also from Theorem

2.2e,f). The bound on $|Y_2^{(r')} - f_2(D_2^{(r)})|$ follows from $Y_2^{(r'-1)} \subset f_2(D_2^{(r)})$, since by that containment the set $Y_2^{(r')} - f_2(D_2^{(r)})$ is a proper subset of the column $Y_2^{r'}$.

9 Appendix 2: Glossary of notation

• $\mathbf{G} = [\mathbf{a_1} \times \mathbf{a_2} \times \cdots \times \mathbf{a_k}]$: the k-dimensional grid, a graph with vertex set $V(G) = \{x = (x_1, x_2, \ldots, x_k) : x_i \text{ an integer, } 1 \leq x_i \leq a_i\}$ and edge set $E(G) = \{xy : \sum_{i=1}^{k} |x_i - y_i| = 1\}$
• $\mathbf{P(t)}$: the path graph on t vertices
• $\mathbf{e_i} = \lceil log_2(a_1 a_2 \cdots a_i) \rceil$ for $1 \leq i \leq k$, with $e_0 = 0$
• **i-page**: a subgraph of G obtained by fixing the last $k - i$ coordinates, and letting the first i coordinates vary over all their possible values
• $\mathbf{D_i^r}$, $\mathbf{D_i^{(r)}}$, $\mathbf{D_i^r(j)}$: D_i^r is the r'th i-page of G under the ordering of i-pages given as follows. Let D_i and D_i' be two i-pages, with fixed last $k - i$ coordinate values $c_{i+1}, c_{i+2}, \ldots, c_k$ and $c'_{i+1}, c'_{i+2}, \ldots, c'_k$ respectively. Then $D_i \prec_i D_i'$ in this ordering if at the maximum index t, $i + 1 \leq t \leq k$, where $c_t \neq c'_t$ we have $c_t < c'_t$. $D_i^{(r)} = \cup_{j=1}^{r} D_i^j$. $D_i^r(j) = D_{i-1}^{(r-1)a_i + j}$, and we regard $D_i^r(j)$ as the j'th $(i-1)$-subpage of D_i^r under the ordering of $(i-1)$-subpages of D_i^r induced by $\prec_{i-1}$.
• $\mathbf{P_i} = a_{i+1} a_{i+2} \ldots a_k$, the number of i-pages in G.
• $\langle \mathbf{Y_i} \rangle$ for $1 \leq i \leq k$: the i-dimensional grid given by
$\langle Y_i \rangle = P(2^{e_1}) \times P(2^{e_2 - e_1}) \times P(2^{e_3 - e_2}) \times \cdots \times P(2^{e_i - e_{i-1}})$. Note that $\langle Y_i \rangle$ is a spanning subgraph of Q_{e_i}, the hypercube of dimension e_i.
• $\mathbf{Y_i}$: the i-dimensional grid given by $Y_i = \langle Y_{i-1} \rangle \times P(l_i)$, where l_i is any integer satisfying $l_i \geq 2^{e_k - e_{i-1}}$
• $\mathbf{(i-1)}$**-level of** $\mathbf{Y_i}$: the subgraph of Y_i consisting of all points with some fixed value c in the i'th coordinate, $1 \leq c \leq l_i$.
• $\mathbf{Y_i^j}$: Y_i^j is the $(i-1)$-level of Y_t consisting of all points with fixed coordinate value j in the i'th coordinate. $(x_1, x_2, \ldots, x_{t-1}, j)$.
• $\mathbf{Y_i^{(r)}} = \bigcup_{j=1}^{r} Y_i^j$.
• **i-section of** $\mathbf{Y_i}$, $\mathbf{S_i^j}$, $\mathbf{S_i^j(G)}$, $\mathbf{S_i^j(G)'}$, $\mathbf{S_i^j(c)}$: $S_i^j = \bigcup_{j=a}^{b} Y_i^j$, where $a = 1 + (j-1)2^{e_i - e_{i-1}}$ and $b = j2^{e_i - e_{i-1}}$. We call S_i^j the j'th "i-section" of Y_i. $S_i^j(G)$ is the set of points of S_i^j lying in nonblank $(i-1)$-levels of S_i^j. $S_i^j(G)' = S_i^j(G) \cap (I_i \circ f_i)(G)$, so $S_i^j(G)'$ is the set of points lying in nonblank columns of S_i^j which are also in the image of $I_i \circ f_i$. $S_i^j(c)$ is the c'th nonblank $(i-1)$-level of S_i^j in order of increasing i'th coordinate. We interpret $S_i^j(c)$ for certain c using wraparound (see the comments preceding Theorem 6.2).
• $\mathbf{S_i^{(r)}}$, $\mathbf{S_i^{(r)}(G)}$, $\mathbf{S_i^{(r)}(G)'}$: $S_i^{(r)} = \bigcup_{j=1}^{r} S_i^j$, the union of the first r many i-sections of Y_i. $S_i^{(r)}(G)$ is the set points in $S_i^{(r)}$ lying in nonblank $(i-1)$-levels of $S_i^{(r)}$. $S_i^{(r)}(G)' = S_i^{(r)}(G) \cap (I_i \circ f_i)(G)$.
• $\mathbf{s_i(j)}$: the number of $(i-1)$-levels in S_i^j which are designated blank. By construction we have $s_i(j) = 2^{e_i - e_{i-1}} - \lceil \frac{a_1 a_2 \cdots a_i}{2^{e_{i-1}}} \rceil + \lfloor j\phi_i \rfloor - \lfloor (j-1)\phi_i \rfloor$, where $\phi_i = \lceil \frac{a_1 a_2 \cdots a_i}{2^{e_{i-1}}} \rceil - \frac{a_1 a_2 \cdots a_i}{2^{e_{i-1}}}$.
• $\mathbf{F(i)}$: the $P_i \times 2^{e_i - e_{i-1}}$, $(0,1)$ matrix $F(i) = f_{uv}(i)$ that encodes which $(i-1)$-levels of $S_i^{(P_i)}$ are to be designated blank as follows. We have $f_{uv}(i) = 1$ if the v'th $(i-1)$-level of S_i^u (which is $Y_i^{v+(u-1)2^{e_i - e_{i-1}}}$) is blank, and $f_{uv}(i) = 0$ otherwise. In particular, the sum of entries in the u'th row of $F(i)$, being the number of blank $(i-1)$-levels in S_i^u, is $s_i(u)$. Note that $F(i)$ is

constructed in the procedure following Corollary 4.5.

• $\mathbf{N_r(d)}$, $\mathbf{N_{i,r}(d)}$: For a fixed i and $1 \le r \le P_i$, $N_r(d)$ is the column index of the d'th 0 from the left of row r of matrix $F(i)$. When i can vary, we let $N_{i,r}(d)$ be the $N_r(d)$ just defined relative to matrix $F(i)$. Further we interpret the function $N_r(*)$ using "wraparound" (see the comments preceding Corollary 4.5).

• $\mathbf{Opt(G)}$, $\mathbf{Opt'(G)}$ for $G = [a_1 \times a_2 \times \cdots \times a_k]$: $Opt(G) = Q_n$ with $n = \lceil log_2(|V(G)|) \rceil$. This is the hypercube of smallest dimension having at least as many vertices as G. $Opt'(G) = P(2^{e_1}) \times P(2^{e_2-e_1}) \times P(2^{e_3-e_2}) \times \cdots \times P(2^{e_k-e_{k-1}})$, a spanning subgraph of $Opt(G)$.

• $\mathbf{u_i = u_i(G)}$: For G a k-dimensional grid, we let $u_i = \lceil \frac{|G|}{2^{e_i-1}} \rceil$. For each i, $2 \le i \le k$, we constructed a map $f_i : G \to Y_i^{(u_i)}$; that is, a map of G into the first u_i many $(i-1)$-levels of Y_i. Each such $(i-1)$-level has size $2^{e_{i-1}}$, so u_i is the minimum number of $(i-1)$-levels required in the image of any one to one map $G \to Y_i$.

• $\mathbf{x_{i \to j}} = (x_i, x_{i+1}, x_{i+2}, \ldots, , x_{j-1}, x_j)$, for a t-tuple $x = (x_1, x_2, x_3, \ldots, x_t)$, $1 \le i, j \le t$.

• $\mathbf{I_i}$: the "inflation" map $I_i : f_i(G) \to S_i^{(P_i)}(G)$ defined in step 3 (The Inflation Step) of the construction of f_{i+1} as follows. For any $z = (z_1, z_2, \ldots, z_i) \in f_i(G)$, let $I_i(z) = (z_1, z_2, \ldots, z_{i-1}, z_i')$, where $Y_i^{z_i'}$ is the z_i'th nonblank level in $S_i^{(P_i)}$ in order of increasing i'th coordinate.

• $\boldsymbol{\sigma_i}$: the "stacking" map $\sigma_i : S_i^{(P_i)}(G)' \to S_i^1 \times P(u_{i+1}) \subseteq Y_{i+1}^{(u_{i+1})}$ defined in step 4 (The Stacking Step) of the construction of f_{i+1}. Informally, σ_i stacks the sets $S_i^r(G)'$ "on top of" $S_i^1 \cong \langle Y_i \rangle$ in order of increasing r as follows. Suppose $x = (x_1, x_2, \ldots, x_i)$ lies in $S_i^r(G)'$. Then let $\sigma_i(x) = (x_1, x_2, \ldots, x_{i-1}, \bar{x}_i, c)$, where $\bar{x}_i$, $1 \le \bar{x}_i \le 2^{e_i - e_{i-1}}$, is the integer congruent to x_i mod $2^{e_i - e_{i-1}}$, and $c = \sigma_i(x)_{i+1}$ is the number of images $\sigma_i(y)$ with $y \in S_i^j(G)$, $1 \le j \le r$, and $\sigma_i(x)_{1 \to i} = \sigma_i(y)_{1 \to i}$.

• $\bar{\boldsymbol{\sigma}}_{\mathbf{i}}$: $\bar{\sigma}_i$ is the projection of σ_i onto the first i coordinates. So for $z \in S_i^{(P_i)}(G)'$, we have $\bar{\sigma}_i(z) = \sigma_i(z)_{1 \to i}$. By Lemma 5.1c, the restriction of $\bar{\sigma}_i$ to any one i-section is one to one.

• $\mathbf{Stack_i(x, r)}$: for $x \in S_{i-1}^1 \cong \langle Y_{i-1} \rangle$ and $1 \le r \le P_{i-1}$ (defined in the paragraph just preceding Lemma 5.1): $Stack_i(x, r) = \{z = \sigma_{i-1}(y) : z_{1 \to i-1} = x \text{ and } y \in S_{i-1}^{(r)}(G)'\}$. We regard $Stack_i(x, r)$ as a stack, with "height" extending into the i'th dimension, addressed by $x \in \langle Y_{i-1} \rangle$. Its elements are images $\sigma_{i-1}(y) \in Y_i$, with $y \in S_{i-1}^{(r)}(G)'$, having projection $\sigma_{i-1}(y)_{1 \to i-1} = x$ onto the first $i-1$ coordinates.

• $\mathbf{[r]_i} = \max \{|Stack_i(x, r)| : x \in \langle Y_{i-1} \rangle\}$. This is the maximum "height" of $Stack_i(x, r)$, over all $x \in \langle Y_{i-1} \rangle$.

• $\mathbf{v_{i,r}(S)}$: for $S \subseteq Y_i^{([r]_i)}$, let $v_{i,r}(S) = |S \cap f_i(D_{i-1}^{(r)})|$.

• $\nu_{\mathbf{i}}(\mathbf{z})$: For $z = (I_i \circ f_i)(x)$, where $x \in D_i^r$, $\nu_i(z)$ is the integer such that $z \in S_i^r(\nu_i(z))$. That is, z belongs to the $\nu_i(z)$'th nonblank $(i-1)$-level (ordered by increasing i-coordinate) of the i-section S_i^r containing z. We interpret differences $\nu_i(z') - \nu_i(z'')$ using wraparound (see the comments preceding Theorem 6.2).

10 Appendix 3: Figures



Figure 1: a) The matrix R, and b) the corresponding map $f : G(5) \to Y_2$



Figure 2: The map $f_2 : [3 \times 7 \times 4 \times a_4] \to Y_2$, with $a_4 > 1$



Figure 3: a) $I_2 : Y_2^{(21)} \to S_2^{(4)}$, used for constructing $f_3([3 \times 7 \times 4])$; a) and b) combined $I_2 : Y_2^{(63)} \to S_2^{(12)}$, used for constructing $f_3([3 \times 7 \times 4 \times 3])$



Figure 4: The stacking map $\sigma_2 : (I_2 \circ f_2)([3 \times 7 \times 4]) \to Y_3^{(3)}$; stacking the four sections S_2^j, $1 \leq j \leq 4$



Figure 5: σ_2 applied to the first 7 and all 12 sections S_2^j, yielding the map $f_3 : [3 \times 7 \times 4 \times 3] \to Y_3^{(8)}$



Figure 6: Contents of stacks in columns 4 and 5 after stacking the first 7 sections S_2^j, $1 \leq j \leq 7$; points are labeled by their preimages in G



Figure 7: The stacking map $\sigma_3(S_3^{(3)}) \subseteq Y_4^{(2)}$, yielding the map $f_4 : [3 \times 7 \times 4 \times 3] \to Y_4^{(2)}$



Figure 8: Three individual stacks, each addressed by points of $S_3^1 \cong \langle Y_3 \rangle$, each of height 2, extending into the 4'th dimension